

# Submodular Functions, Optimization, and Applications to Machine Learning

— Spring Quarter, Lecture 4 —

[http://j.ee.washington.edu/~bilmes/classes/ee596b\\_spring\\_2014/](http://j.ee.washington.edu/~bilmes/classes/ee596b_spring_2014/)

Prof. Jeff Bilmes


University of Washington, Seattle  
Department of Electrical Engineering

<http://melodi.ee.washington.edu/~bilmes>

April 9th, 2014



$$f(A) + f(B) \geq f(A \cup B) + f(A \cap B)$$



# Cumulative Outstanding Reading

- Read chapter 1 from Fujishige's book.

# Announcements, Assignments, and Reminders

- our room (Mueller Hall Room 154) is changed!
- Please do use our discussion board ([https://canvas.uw.edu/courses/895956/discussion\\_topics](https://canvas.uw.edu/courses/895956/discussion_topics)) for all questions, comments, so that all will benefit from them being answered.
- Weekly Office Hours: Wednesdays, 5:00-5:50, or by skype or google hangout (email me).

# Class Road Map - IT-I

- L1 (3/31): Motivation, Applications, & Basic Definitions
- L2: (4/2): Applications, Basic Definitions, Properties
- L3: More examples and properties (e.g., closure properties), and examples, spanning trees
- L4: proofs of equivalent definitions, independence, start matroids
- L5:
- L6:
- L7:
- L8:
- L9:
- L10:
- L11:
- L12:
- L13:
- L14:
- L15:
- L16:
- L17:
- L18:
- L19:
- L20:

Finals Week: June 9th-13th, 2014.



# Summary so far

- Summing: if  $\alpha_i \geq 0$  and  $f_i : 2^V \rightarrow \mathbb{R}$  is submodular, then so is  $\sum_i \alpha_i f_i$ .
- Restrictions:  $f'(A) = f(A \cap S)$
- max:  $f(A) = \max_{j \in A} c_j$  and facility location.
- Log determinant  $f(A) = \log \det(\Sigma_A)$

# Concave over non-negative modular

Let  $m \in \mathbb{R}_+^E$  be a modular function, and  $g$  a concave function over  $\mathbb{R}$ . Define  $f : 2^E \rightarrow \mathbb{R}$  as

$$f(A) = g(m(A)) \quad (4.35)$$

then  $f$  is submodular.

**Proof.**

Given  $A \subseteq B \subseteq E \setminus v$ , we have  $0 \leq a = m(A) \leq b = m(B)$ , and  $0 \leq c = m(v)$ . For  $g$  concave, we have  $g(a + c) - g(a) \geq g(b + c) - g(b)$ , and thus

$$g(m(A) + m(v)) - g(m(A)) \geq g(m(B) + m(v)) - g(m(B)) \quad (4.36)$$



A form of converse is true as well.

# Concave composed with non-negative modular

## Theorem 4.2.1

*Given a ground set  $V$ . The following two are equivalent:*

- ① *For all modular functions  $m : 2^V \rightarrow \mathbb{R}_+$ , then  $f : 2^V \rightarrow \mathbb{R}$  defined as  $f(A) = g(m(A))$  is submodular*
- ②  *$g : \mathbb{R}_+ \rightarrow \mathbb{R}$  is concave.*

- If  $g$  is non-decreasing concave, then  $f$  is polymatroidal.
- Sums of concave over modular functions are submodular

$$f(A) = \sum_{i=1}^K g_i(m_i(A)) \quad (4.35)$$

- Very large class of functions, including graph cut, bipartite neighborhoods, set cover (Stobbe & Krause).
- However, Vondrak showed that a graphic matroid rank function over  $K_4$  (we'll define this after we define matroids) are not members.

# Composition of non-decreasing submodular and non-decreasing concave

## Theorem 4.2.1

*Given two functions, one defined on sets*

$$f : 2^V \rightarrow \mathbb{R} \quad (4.35)$$

*and another continuous valued one:*

$$g : \mathbb{R} \rightarrow \mathbb{R} \quad (4.36)$$

*the composition formed as  $h = g \circ f : 2^V \rightarrow \mathbb{R}$  (defined as  $h(S) = g(f(S))$ ) is nondecreasing submodular, if  $g$  is non-decreasing concave and  $f$  is nondecreasing submodular.*

# Monotone difference of two functions

Let  $f$  and  $g$  both be submodular functions on subsets of  $V$  and let  $(f - g)(\cdot)$  be either monotone increasing or monotone decreasing. Then  $h : 2^V \rightarrow R$  defined by

$$h(A) = \min(f(A), g(A)) \quad (4.35)$$

is submodular.

Proof.

If  $h(A)$  agrees with either  $f$  or  $g$  on **both**  $X$  and  $Y$ , and since

$$f(X) + f(Y) \geq f(X \cup Y) + f(X \cap Y) \quad (4.36)$$

$$g(X) + g(Y) \geq g(X \cup Y) + g(X \cap Y), \quad (4.37)$$

the result (Equation ??) follows since

$$\begin{aligned} f(X) + f(Y) &\geq \min(f(X \cup Y), g(X \cup Y)) + \min(f(X \cap Y), g(X \cap Y)) \\ g(X) + g(Y) &\geq \min(f(X \cup Y), g(X \cup Y)) + \min(f(X \cap Y), g(X \cap Y)) \end{aligned} \quad (4.38)$$

# Saturation via the $\min(\cdot)$ function

Let  $f : 2^V \rightarrow \mathbb{R}$  be an monotone increasing or decreasing submodular function and let  $k$  be a constant. Then the function  $h : 2^V \rightarrow \mathbb{R}$  defined by

$$h(A) = \min(k, f(A)) \quad (4.37)$$

is submodular.

Proof.

For constant  $k$ , we have that  $(f - k)$  is increasing (or decreasing) so this follows from the previous result. □

Note also,  $g(a) = \min(k, a)$  for constant  $k$  is a non-decreasing concave function, so when  $f$  is monotone nondecreasing submodular, we can use the earlier result about composing a monotone concave function with a monotone submodular function to get a version of this.

# Gain Notation

It will also be useful to extend this to sets.

Let  $A, B$  be any two sets. Then

$$f(A|B) \triangleq f(A \cup B) - f(B) \quad (4.41)$$

So when  $j$  is any singleton

$$f(j|B) = f(\{j\}|B) = f(\{j\} \cup B) - f(B) \quad (4.42)$$

Note that this is inspired from information theory and the notation used for conditional entropy  $H(X_A|X_B) = H(X_A, X_B) - H(X_B)$ .

# Other properties

- Any submodular function  $h : 2^V \rightarrow \mathbb{R}$  can be represented as the difference between two submodular functions, i.e.,  
 $h(A) = f(A) - g(A)$  where both  $f$  and  $g$  are submodular.



# Other properties

- Any submodular function  $h : 2^V \rightarrow \mathbb{R}$  can be represented as the difference between two submodular functions, i.e.,  $h(A) = f(A) - g(A)$  where both  $f$  and  $g$  are submodular.
- Any submodular function  $f$  can be represented as a sum of a normalized monotone non-decreasing submodular function and a modular function,  $f = \bar{f} + m$

# Other properties

- Any submodular function  $h : 2^V \rightarrow \mathbb{R}$  can be represented as the difference between two submodular functions, i.e.,  
 $h(A) = f(A) - g(A)$  where both  $f$  and  $g$  are submodular.
- Any submodular function  $f$  can be represented as a sum of a normalized monotone non-decreasing submodular function and a modular function,  $f = \bar{f} + m$
- Any function  $h$  can be represented as the difference between two monotone non-decreasing submodular functions.

# Submodular Definitions

## Definition 4.3.2 (submodular concave)

A function  $f : 2^V \rightarrow \mathbb{R}$  is **submodular** if for any  $A, B \subseteq V$ , we have that:

$$f(A) + f(B) \geq f(A \cup B) + f(A \cap B) \quad (4.2)$$

An alternate and (as we will soon see) equivalent definition is:

## Definition 4.3.3 (diminishing returns)

A function  $f : 2^V \rightarrow \mathbb{R}$  is **submodular** if for any  $A \subseteq B \subset V$ , and  $v \in V \setminus B$ , we have that:

$$f(A \cup \{v\}) - f(A) \geq f(B \cup \{v\}) - f(B) \quad (4.3)$$

This means that the incremental “value”, “gain”, or “cost” of  $v$  decreases (diminishes) as the context in which  $v$  is considered grows from  $A$  to  $B$ .

# Submodular Definition: Group Diminishing Returns

An alternate and equivalent definition is:

## Definition 4.3.1 (group diminishing returns)

A function  $f : 2^V \rightarrow \mathbb{R}$  is submodular if for any  $A \subseteq B \subset V$ , and  $C \subseteq V \setminus B$ , we have that:

$$f(A \cup C) - f(A) \geq f(B \cup C) - f(B) \quad (4.1)$$

This means that the incremental “value” or “gain” of **set**  $C$  decreases as the context in which  $C$  is considered grows from  $A$  to  $B$  (diminishing returns)

# Submodular Definition Basic Equivalencies

We want to show that **Submodular Concave** (Definition 4.3.2), **Diminishing Returns** (Definition 4.3.3), and **Group Diminishing Returns** (Definition 4.3.1) are identical.

# Submodular Definition Basic Equivalencies

We want to show that **Submodular Concave** (Definition 4.3.2), **Diminishing Returns** (Definition 4.3.3), and **Group Diminishing Returns** (Definition 4.3.1) are identical. We will show that:

- Submodular Concave  $\Rightarrow$  Diminishing Returns
- Diminishing Returns  $\Rightarrow$  Group Diminishing Returns
- Group Diminishing Returns  $\Rightarrow$  Submodular Concave

# Submodular Concave $\Rightarrow$ Diminishing Returns

$$f(S) + f(T) \geq f(S \cup T) + f(S \cap T) \Rightarrow f(v|A) \geq f(v|B), A \subseteq B \subseteq V \setminus v.$$

- Assume Submodular concave, so  $\forall S, T$  we have  $f(S) + f(T) \geq f(S \cup T) + f(S \cap T)$ .



# Submodular Concave $\Rightarrow$ Diminishing Returns

$$f(S) + f(T) \geq f(S \cup T) + f(S \cap T) \Rightarrow f(v|A) \geq f(v|B), A \subseteq B \subseteq V \setminus v.$$

- Assume Submodular concave, so  $\forall S, T$  we have  $f(S) + f(T) \geq f(S \cup T) + f(S \cap T)$ .
- Given  $A, B$  and  $v \in V$  such that:  $A \subseteq B \subseteq V \setminus \{v\}$ , we have from submodular concave that:

$$f(A + v) + f(B) \geq f(B + v) + f(A) \quad (4.2)$$





# Submodular Concave $\Rightarrow$ Diminishing Returns

$$f(S) + f(T) \geq f(S \cup T) + f(S \cap T) \Rightarrow f(v|A) \geq f(v|B), A \subseteq B \subseteq V \setminus v.$$

- Assume Submodular concave, so  $\forall S, T$  we have  $f(S) + f(T) \geq f(S \cup T) + f(S \cap T)$ .
- Given  $A, B$  and  $v \in V$  such that:  $A \subseteq B \subseteq V \setminus \{v\}$ , we have from submodular concave that:

$$f(A + v) + f(B) \geq f(B + v) + f(A) \quad (4.2)$$

- Rearranging, we have

$$f(A + v) - f(A) \geq f(B + v) - f(B) \quad (4.3)$$



# Diminishing Returns $\Rightarrow$ Group Diminishing Returns

$$f(v|S) \geq f(v|T), S \subseteq T \subseteq V \setminus v \Rightarrow f(C|A) \geq f(C|B), A \subseteq B \subseteq V \setminus C.$$

Let  $C = \{c_1, c_2, \dots, c_k\}$ . Then **diminishing returns** implies

$$f(A \cup C) - f(A) \tag{4.4}$$

$$= f(A \cup C) - \sum_{i=1}^{k-1} \left( f(A \cup \{c_1, \dots, c_i\}) - f(A \cup \{c_1, \dots, c_{i-1}\}) \right) - f(A) \tag{4.5}$$

$$= \sum_{i=1}^k \left( f(A \cup \{c_1 \dots c_i\}) - f(A \cup \{c_1 \dots c_{i-1}\}) \right) \tag{4.6}$$

$$\geq \sum_{i=1}^k \left( f(B \cup \{c_1 \dots c_i\}) - f(B \cup \{c_1 \dots c_{i-1}\}) \right) \tag{4.7}$$

$$= f(B \cup C) - \sum_{i=1}^{k-1} \left( f(B \cup \{c_1, \dots, c_i\}) - f(B \cup \{c_1, \dots, c_{i-1}\}) \right) - f(B) \tag{4.8}$$

$$= f(B \cup C) - f(B) \tag{4.9}$$



# Group Diminishing Returns $\Rightarrow$ Submodular Concave

$$f(U|S) \geq f(U|T), S \subseteq T \subseteq V \setminus U \Rightarrow f(A) + f(B) \geq f(A \cup B) + f(A \cap B).$$

Assume **group diminishing returns**. Assume  $A \neq B$  otherwise trivial.

Define  $A' = A \cap B$ ,  $C = A \setminus B$ , and  $B' = B$ . Then since  $A' \subseteq B'$ ,

$$f(A' + C) - f(A') \geq f(B' + C) - f(B') \quad (4.10)$$

giving

$$f(A' + C) + f(B') \geq f(B' + C) + f(A') \quad (4.11)$$

or

$$f(A \cap B + A \setminus B) + f(B) \geq f(B + A \setminus B) + f(A \cap B) \quad (4.12)$$

which is the same as the submodular concave condition

$$f(A) + f(B) \geq f(A \cup B) + f(A \cap B) \quad (4.13)$$

## Submodular Definition: Four Points

### Definition 4.3.2 (“singleton”, or “four points”)

A function  $f : 2^V \rightarrow \mathbb{R}$  is submodular iff for any  $A \subset V$ , and any  $a, b \in V \setminus A$ , we have that:

$$f(A \cup \{a\}) + f(A \cup \{b\}) \geq f(A \cup \{a, b\}) + f(A) \quad (4.14)$$

# Submodular Definition: Four Points

## Definition 4.3.2 (“singleton”, or “four points”)

A function  $f : 2^V \rightarrow \mathbb{R}$  is submodular iff for any  $A \subset V$ , and any  $a, b \in V \setminus A$ , we have that:

$$f(A \cup \{a\}) + f(A \cup \{b\}) \geq f(A \cup \{a, b\}) + f(A) \quad (4.14)$$

This follows immediately from **diminishing returns**.

# Submodular Definition: Four Points

## Definition 4.3.2 (“singleton”, or “four points”)

A function  $f : 2^V \rightarrow \mathbb{R}$  is submodular iff for any  $A \subset V$ , and any  $a, b \in V \setminus A$ , we have that:

$$f(A \cup \{a\}) + f(A \cup \{b\}) \geq f(A \cup \{a, b\}) + f(A) \quad (4.14)$$

This follows immediately from **diminishing returns**. To achieve **diminishing returns**, assume  $A \subset B$  with  $B \setminus A = \{b_1, b_2, \dots, b_k\}$ . Then

$$f(A + a) - f(A) \geq f(A + b_1 + a) - f(A + b_1) \quad (4.15)$$

$$\geq f(A + b_1 + b_2 + a) - f(A + b_1 + b_2) \quad (4.16)$$

$$\geq \dots \quad (4.17)$$

$$\geq f(A + b_1 + \dots + b_k + a) - f(A + b_1 + \dots + b_k) \quad (4.18)$$

$$= f(B + a) - f(B) \quad (4.19)$$

# Submodular Definitions

## Theorem 4.3.3

Given function  $f : 2^V \rightarrow \mathbb{R}$ , then

$$f(A) + f(B) \geq f(A \cup B) + f(A \cap B) \text{ for all } A, B \subseteq V \quad (\text{SC})$$

if and only if

$$f(v|X) \geq f(v|Y) \text{ for all } X \subseteq Y \subseteq V \text{ and } v \notin B \quad (\text{DR})$$

# Submodular Definitions

## Theorem 4.3.3

Given function  $f : 2^V \rightarrow \mathbb{R}$ , then

$$f(A) + f(B) \geq f(A \cup B) + f(A \cap B) \text{ for all } A, B \subseteq V \quad (\text{SC})$$

if and only if

$$f(v|X) \geq f(v|Y) \text{ for all } X \subseteq Y \subseteq V \text{ and } v \notin B \quad (\text{DR})$$

## Proof.

(SC) $\Rightarrow$ (DR): Set  $A \leftarrow X \cup \{v\}$ ,  $B \leftarrow Y$ . Then  $A \cup B = B \cup \{v\}$  and  $A \cap B = X$  and  $f(A) - f(A \cap B) \geq f(A \cup B) - f(B)$  implies (DR).

(DR) $\Rightarrow$ (SC): Order  $A \setminus B = \{v_1, v_2, \dots, v_r\}$  arbitrarily. Then  $f(v_i|A \cap B \cup \{v_1, v_2, \dots, v_{i-1}\}) \geq f(v_i|B \cup \{v_1, v_2, \dots, v_{i-1}\})$ ,  $i \in [r - 1]$

Applying telescoping summation to both sides, we get:

$$\sum_{i=0}^r f(v_i|A \cap B \cup \{v_1, v_2, \dots, v_{i-1}\}) \geq \sum_{i=0}^r f(v_i|B \cup \{v_1, v_2, \dots, v_{i-1}\})$$

or

$$f(A) - f(A \cap B) \geq f(A \cup B) - f(B)$$



# Use of gain: submodular bounds of a difference

- Given submodular  $f$ , and given you have  $C, D \subseteq E$  with either  $D \supseteq C$  or  $D \subseteq C$ , and have an expression of the form:

$$f(C) - f(D) \tag{4.20}$$

# Use of gain: submodular bounds of a difference

- Given submodular  $f$ , and given you have  $C, D \subseteq E$  with either  $D \supseteq C$  or  $D \subseteq C$ , and have an expression of the form:

$$f(C) - f(D) \tag{4.20}$$

- If  $D \supseteq C$ , then for any  $X$  with  $D = C \cup X$  then

$$f(C) - f(D) = f(C) - f(C \cup X) \geq f(C \cap X) - f(X)$$

$$\tag{4.22}$$

# Use of gain: submodular bounds of a difference

- Given submodular  $f$ , and given you have  $C, D \subseteq E$  with either  $D \supseteq C$  or  $D \subseteq C$ , and have an expression of the form:

$$f(C) - f(D) \tag{4.20}$$

- If  $D \supseteq C$ , then for any  $X$  with  $D = C \cup X$  then

$$f(C) - f(D) = f(C) - f(C \cup X) \geq f(C \cap X) - f(X) \tag{4.21}$$

or

$$f(C \cup X|C) \leq f(X|C \cap X) \tag{4.22}$$

# Use of gain: submodular bounds of a difference

- Given submodular  $f$ , and given you have  $C, D \subseteq E$  with either  $D \supseteq C$  or  $D \subseteq C$ , and have an expression of the form:

$$f(C) - f(D) \tag{4.20}$$

- If  $D \supseteq C$ , then for any  $X$  with  $D = C \cup X$  then

$$f(C) - f(D) = f(C) - f(C \cup X) \geq f(C \cap X) - f(X) \tag{4.21}$$

or

$$f(C \cup X|C) \leq f(X|C \cap X) \tag{4.22}$$

- Alternatively, if  $D \subseteq C$ , given any  $Y$  such that  $D = C \cap Y$  then

$$f(C) - f(D) = f(C) - f(C \cap Y) \geq f(C \cup Y) - f(Y)$$

$$\tag{4.24}$$

# Use of gain: submodular bounds of a difference

- Given submodular  $f$ , and given you have  $C, D \subseteq E$  with either  $D \supseteq C$  or  $D \subseteq C$ , and have an expression of the form:

$$f(C) - f(D) \tag{4.20}$$

- If  $D \supseteq C$ , then for any  $X$  with  $D = C \cup X$  then

$$f(C) - f(D) = f(C) - f(C \cup X) \geq f(C \cap X) - f(X) \tag{4.21}$$

or

$$f(C \cup X|C) \leq f(X|C \cap X) \tag{4.22}$$

- Alternatively, if  $D \subseteq C$ , given any  $Y$  such that  $D = C \cap Y$  then

$$f(C) - f(D) = f(C) - f(C \cap Y) \geq f(C \cup Y) - f(Y) \tag{4.23}$$

or

$$f(C|C \cap Y) \geq f(C \cup Y|Y) \tag{4.24}$$

# Use of gain: submodular bounds of a difference

- Given submodular  $f$ , and given you have  $C, D \subseteq E$  with either  $D \supseteq C$  or  $D \subseteq C$ , and have an expression of the form:

$$f(C) - f(D) \tag{4.20}$$

- If  $D \supseteq C$ , then for any  $X$  with  $D = C \cup X$  then

$$f(C) - f(D) = f(C) - f(C \cup X) \geq f(C \cap X) - f(X) \tag{4.21}$$

or

$$f(C \cup X|C) \leq f(X|C \cap X) \tag{4.22}$$

- Alternatively, if  $D \subseteq C$ , given any  $Y$  such that  $D = C \cap Y$  then

$$f(C) - f(D) = f(C) - f(C \cap Y) \geq f(C \cup Y) - f(Y) \tag{4.23}$$

or

$$f(C|C \cap Y) \geq f(C \cup Y|Y) \tag{4.24}$$

- Equations (4.22) and (4.24) have same form.

# Many (Equivalent) Definitions of Submodularity

$$f(A) + f(B) \geq f(A \cup B) + f(A \cap B), \quad \forall A, B \subseteq V \quad (4.25)$$

# Many (Equivalent) Definitions of Submodularity

$$f(A) + f(B) \geq f(A \cup B) + f(A \cap B), \quad \forall A, B \subseteq V \quad (4.25)$$

$$f(j|S) \geq f(j|T), \quad \forall S \subseteq T \subseteq V, \text{ with } j \in V \setminus T \quad (4.26)$$



# Many (Equivalent) Definitions of Submodularity

$$f(A) + f(B) \geq f(A \cup B) + f(A \cap B), \quad \forall A, B \subseteq V \quad (4.25)$$

$$f(j|S) \geq f(j|T), \quad \forall S \subseteq T \subseteq V, \text{ with } j \in V \setminus T \quad (4.26)$$

$$f(C|S) \geq f(C|T), \forall S \subseteq T \subseteq V, \text{ with } C \subseteq V \setminus T \quad (4.27)$$

# Many (Equivalent) Definitions of Submodularity

$$f(A) + f(B) \geq f(A \cup B) + f(A \cap B), \quad \forall A, B \subseteq V \quad (4.25)$$

$$f(j|S) \geq f(j|T), \quad \forall S \subseteq T \subseteq V, \text{ with } j \in V \setminus T \quad (4.26)$$

$$f(C|S) \geq f(C|T), \quad \forall S \subseteq T \subseteq V, \text{ with } C \subseteq V \setminus T \quad (4.27)$$

$$f(j|S) \geq f(j|S \cup \{k\}), \quad \forall S \subseteq V \text{ with } j \in V \setminus (S \cup \{k\}) \quad (4.28)$$

# Many (Equivalent) Definitions of Submodularity

$$f(A) + f(B) \geq f(A \cup B) + f(A \cap B), \quad \forall A, B \subseteq V \quad (4.25)$$

$$f(j|S) \geq f(j|T), \quad \forall S \subseteq T \subseteq V, \text{ with } j \in V \setminus T \quad (4.26)$$

$$f(C|S) \geq f(C|T), \quad \forall S \subseteq T \subseteq V, \text{ with } C \subseteq V \setminus T \quad (4.27)$$

$$f(j|S) \geq f(j|S \cup \{k\}), \quad \forall S \subseteq V \text{ with } j \in V \setminus (S \cup \{k\}) \quad (4.28)$$

$$f(A \cup B|A \cap B) \leq f(A|A \cap B) + f(B|A \cap B), \quad \forall A, B \subseteq V \quad (4.29)$$

# Many (Equivalent) Definitions of Submodularity

$$f(A) + f(B) \geq f(A \cup B) + f(A \cap B), \quad \forall A, B \subseteq V \quad (4.25)$$

$$f(j|S) \geq f(j|T), \quad \forall S \subseteq T \subseteq V, \text{ with } j \in V \setminus T \quad (4.26)$$

$$f(C|S) \geq f(C|T), \quad \forall S \subseteq T \subseteq V, \text{ with } C \subseteq V \setminus T \quad (4.27)$$

$$f(j|S) \geq f(j|S \cup \{k\}), \quad \forall S \subseteq V \text{ with } j \in V \setminus (S \cup \{k\}) \quad (4.28)$$

$$f(A \cup B|A \cap B) \leq f(A|A \cap B) + f(B|A \cap B), \quad \forall A, B \subseteq V \quad (4.29)$$

$$f(T) \leq f(S) + \sum_{j \in T \setminus S} f(j|S) - \sum_{j \in S \setminus T} f(j|S \cup T - \{j\}), \quad \forall S, T \subseteq V \quad (4.30)$$

# Many (Equivalent) Definitions of Submodularity

$$f(A) + f(B) \geq f(A \cup B) + f(A \cap B), \quad \forall A, B \subseteq V \quad (4.25)$$

$$f(j|S) \geq f(j|T), \quad \forall S \subseteq T \subseteq V, \text{ with } j \in V \setminus T \quad (4.26)$$

$$f(C|S) \geq f(C|T), \quad \forall S \subseteq T \subseteq V, \text{ with } C \subseteq V \setminus T \quad (4.27)$$

$$f(j|S) \geq f(j|S \cup \{k\}), \quad \forall S \subseteq V \text{ with } j \in V \setminus (S \cup \{k\}) \quad (4.28)$$

$$f(A \cup B|A \cap B) \leq f(A|A \cap B) + f(B|A \cap B), \quad \forall A, B \subseteq V \quad (4.29)$$

$$f(T) \leq f(S) + \sum_{j \in T \setminus S} f(j|S) - \sum_{j \in S \setminus T} f(j|S \cup T - \{j\}), \quad \forall S, T \subseteq V \quad (4.30)$$

$$f(T) \leq f(S) + \sum_{j \in T \setminus S} f(j|S), \quad \forall S \subseteq T \subseteq V \quad (4.31)$$

# Many (Equivalent) Definitions of Submodularity

$$f(A) + f(B) \geq f(A \cup B) + f(A \cap B), \quad \forall A, B \subseteq V \quad (4.25)$$

$$f(j|S) \geq f(j|T), \quad \forall S \subseteq T \subseteq V, \text{ with } j \in V \setminus T \quad (4.26)$$

$$f(C|S) \geq f(C|T), \quad \forall S \subseteq T \subseteq V, \text{ with } C \subseteq V \setminus T \quad (4.27)$$

$$f(j|S) \geq f(j|S \cup \{k\}), \quad \forall S \subseteq V \text{ with } j \in V \setminus (S \cup \{k\}) \quad (4.28)$$

$$f(A \cup B|A \cap B) \leq f(A|A \cap B) + f(B|A \cap B), \quad \forall A, B \subseteq V \quad (4.29)$$

$$f(T) \leq f(S) + \sum_{j \in T \setminus S} f(j|S) - \sum_{j \in S \setminus T} f(j|S \cup T - \{j\}), \quad \forall S, T \subseteq V \quad (4.30)$$

$$f(T) \leq f(S) + \sum_{j \in T \setminus S} f(j|S), \quad \forall S \subseteq T \subseteq V \quad (4.31)$$

$$f(T) \leq f(S) - \sum_{j \in S \setminus T} f(j|S \setminus \{j\}) + \sum_{j \in T \setminus S} f(j|S \cap T) \quad \forall S, T \subseteq V \quad (4.32)$$

# Many (Equivalent) Definitions of Submodularity

$$f(A) + f(B) \geq f(A \cup B) + f(A \cap B), \quad \forall A, B \subseteq V \quad (4.25)$$

$$f(j|S) \geq f(j|T), \quad \forall S \subseteq T \subseteq V, \text{ with } j \in V \setminus T \quad (4.26)$$

$$f(C|S) \geq f(C|T), \quad \forall S \subseteq T \subseteq V, \text{ with } C \subseteq V \setminus T \quad (4.27)$$

$$f(j|S) \geq f(j|S \cup \{k\}), \quad \forall S \subseteq V \text{ with } j \in V \setminus (S \cup \{k\}) \quad (4.28)$$

$$f(A \cup B|A \cap B) \leq f(A|A \cap B) + f(B|A \cap B), \quad \forall A, B \subseteq V \quad (4.29)$$

$$f(T) \leq f(S) + \sum_{j \in T \setminus S} f(j|S) - \sum_{j \in S \setminus T} f(j|S \cup T - \{j\}), \quad \forall S, T \subseteq V \quad (4.30)$$

$$f(T) \leq f(S) + \sum_{j \in T \setminus S} f(j|S), \quad \forall S \subseteq T \subseteq V \quad (4.31)$$

$$f(T) \leq f(S) - \sum_{j \in S \setminus T} f(j|S \setminus \{j\}) + \sum_{j \in T \setminus S} f(j|S \cap T) \quad \forall S, T \subseteq V \quad (4.32)$$

$$f(T) \leq f(S) - \sum_{j \in S \setminus T} f(j|S \setminus \{j\}), \quad \forall T \subseteq S \subseteq V \quad (4.33)$$

# Equivalent Definitions of Submodularity

We've already seen that  $\text{Eq. 4.25} \equiv \text{Eq. 4.26} \equiv \text{Eq. 4.27} \equiv \text{Eq. 4.28} \equiv \text{Eq. 4.29}$ .



# Equivalent Definitions of Submodularity

We've already seen that  $\text{Eq. 4.25} \equiv \text{Eq. 4.26} \equiv \text{Eq. 4.27} \equiv \text{Eq. 4.28} \equiv \text{Eq. 4.29}$ .

We next show that  $\text{Eq. 4.28} \Rightarrow \text{Eq. 4.30} \Rightarrow \text{Eq. 4.31} \Rightarrow \text{Eq. 4.28}$ .

# Approach

To show these next results, we essentially first use:

$$f(S \cup T) = f(S) + f(T|S) \leq f(S) + \text{upper-bound} \quad (4.34)$$

and

$$f(T) + \text{lower-bound} \leq f(T) + f(S|T) = f(S \cup T) \quad (4.35)$$

# Approach

To show these next results, we essentially first use:

$$f(S \cup T) = f(S) + f(T|S) \leq f(S) + \text{upper-bound} \quad (4.34)$$

and

$$f(T) + \text{lower-bound} \leq f(T) + f(S|T) = f(S \cup T) \quad (4.35)$$

leading to

$$f(T) + \text{lower-bound} \leq f(S) + \text{upper-bound} \quad (4.36)$$

or

$$f(T) \leq f(S) + \text{upper-bound} - \text{lower-bound} \quad (4.37)$$

Eq. 4.28  $\Rightarrow$  Eq. 4.30

Let  $T \setminus S = \{j_1, \dots, j_r\}$  and  $S \setminus T = \{k_1, \dots, k_q\}$ .

First, we upper bound the gain of  $T$  in the context of  $S$ :

$$f(S \cup T) - f(S) = \sum_{t=1}^r \left( f(S \cup \{j_1, \dots, j_t\}) - f(S \cup \{j_1, \dots, j_{t-1}\}) \right) \quad (4.38)$$

$$= \sum_{t=1}^r f(j_t | S \cup \{j_1, \dots, j_{t-1}\}) \leq \sum_{t=1}^r f(j_t | S) \quad (4.39)$$

$$= \sum_{j \in T \setminus S} f(j | S) \quad (4.40)$$

or

$$f(T | S) \leq \sum_{j \in T \setminus S} f(j | S) \quad (4.41)$$

# Eq. 4.28 $\Rightarrow$ Eq. 4.30

Let  $T \setminus S = \{j_1, \dots, j_r\}$  and  $S \setminus T = \{k_1, \dots, k_q\}$ .

Next, lower bound  $S$  in the context of  $T$ :

$$f(S \cup T) - f(T) = \sum_{t=1}^q [f(T \cup \{k_1, \dots, k_t\}) - f(T \cup \{k_1, \dots, k_{t-1}\})] \quad (4.42)$$

$$= \sum_{t=1}^q f(k_t | T \cup \{k_1, \dots, k_t\} \setminus \{k_t\}) \geq \sum_{t=1}^q f(k_t | T \cup S \setminus \{k_t\}) \quad (4.43)$$

$$= \sum_{j \in S \setminus T} f(j | S \cup T \setminus \{j\}) \quad (4.44)$$

# Eq. 4.28 $\Rightarrow$ Eq. 4.30

Let  $T \setminus S = \{j_1, \dots, j_r\}$  and  $S \setminus T = \{k_1, \dots, k_q\}$ .

So we have the upper bound

$$f(T|S) = f(S \cup T) - f(S) \leq \sum_{j \in T \setminus S} f(j|S) \quad (4.45)$$

and the lower bound

$$f(S|T) = f(S \cup T) - f(T) \geq \sum_{j \in S \setminus T} f(j|S \cup T \setminus \{j\}) \quad (4.46)$$

This gives upper and lower bounds of the form

$$f(T) + \text{lower bound} \leq f(S \cup T) \leq f(S) + \text{upper bound}, \quad (4.47)$$

and combining directly the left and right hand side gives the desired inequality.

Eq. 4.30  $\Rightarrow$  Eq. 4.31

This follows immediately since if  $S \subseteq T$ , then  $S \setminus T = \emptyset$ , and the last term of Eq. 4.30 vanishes.

Eq. 4.31  $\Rightarrow$  Eq. 4.28

Here, we set  $T = S \cup \{j, k\}$ ,  $j \notin S \cup \{k\}$  into Eq. 4.31 to obtain

$$f(S \cup \{j, k\}) \leq f(S) + f(j|S) + f(k|S) \quad (4.48)$$

$$= f(S) + f(S + \{j\}) - f(S) + f(S + \{k\}) - f(S) \quad (4.49)$$

$$= f(S + \{j\}) + f(S + \{k\}) - f(S) \quad (4.50)$$

$$= f(j|S) + f(S + \{k\}) \quad (4.51)$$

giving

$$f(j|S \cup \{k\}) = f(S \cup \{j, k\}) - f(S \cup \{k\}) \quad (4.52)$$

$$\leq f(j|S) \quad (4.53)$$



# Submodular Concave

- Why do we call the  $f(A) + f(B) \geq f(A \cup B) + f(A \cap B)$  definition of submodularity, submodular **concave**?

# Submodular Concave

- Why do we call the  $f(A) + f(B) \geq f(A \cup B) + f(A \cap B)$  definition of submodularity, submodular **concave**?
- A continuous twice differentiable function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is concave iff  $\nabla^2 f \preceq 0$  (the Hessian matrix is nonpositive definite).

# Submodular Concave

- Why do we call the  $f(A) + f(B) \geq f(A \cup B) + f(A \cap B)$  definition of submodularity, submodular **concave**?
- A continuous twice differentiable function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is concave iff  $\nabla^2 f \preceq 0$  (the Hessian matrix is nonpositive definite).
- Define a “discrete derivative” or difference operator defined on discrete functions  $f : 2^V \rightarrow \mathbb{R}$  as follows:

$$(\nabla_B f)(A) \triangleq f(A \cup B) - f(A \setminus B) = f(B|(A \setminus B)) \quad (4.54)$$

read as: the derivative of  $f$  at  $A$  in the direction  $B$ .

# Submodular Concave

- Why do we call the  $f(A) + f(B) \geq f(A \cup B) + f(A \cap B)$  definition of submodularity, submodular **concave**?
- A continuous twice differentiable function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is concave iff  $\nabla^2 f \preceq 0$  (the Hessian matrix is nonpositive definite).
- Define a “discrete derivative” or difference operator defined on discrete functions  $f : 2^V \rightarrow \mathbb{R}$  as follows:

$$(\nabla_B f)(A) \triangleq f(A \cup B) - f(A \setminus B) = f(B|(A \setminus B)) \quad (4.54)$$

read as: the derivative of  $f$  at  $A$  in the direction  $B$ .

- Hence, if  $A \cap B = \emptyset$ , then  $(\nabla_B f)(A) = f(B|A)$ .

# Submodular Concave

- Why do we call the  $f(A) + f(B) \geq f(A \cup B) + f(A \cap B)$  definition of submodularity, submodular **concave**?
- A continuous twice differentiable function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is concave iff  $\nabla^2 f \preceq 0$  (the Hessian matrix is nonpositive definite).
- Define a “discrete derivative” or difference operator defined on discrete functions  $f : 2^V \rightarrow \mathbb{R}$  as follows:

$$(\nabla_B f)(A) \triangleq f(A \cup B) - f(A \setminus B) = f(B|(A \setminus B)) \quad (4.54)$$

read as: the derivative of  $f$  at  $A$  in the direction  $B$ .

- Hence, if  $A \cap B = \emptyset$ , then  $(\nabla_B f)(A) = f(B|A)$ .
- Consider a form of second derivative or 2nd difference:

$$\begin{aligned} (\nabla_C \nabla_B f)(A) &= \nabla_C[f(A \cup B) - f(A \setminus B)] \\ &= f(A \cup B \cup C) - f((A \cup C) \setminus B) \\ &\quad - f((A \setminus C) \cup B) + f((A \setminus C) \setminus B) \end{aligned} \quad (4.55)$$

# Submodular Concave

- If the second difference operator everywhere nonpositive:

$$\begin{aligned}
 &f(A \cup B \cup C) - f((A \cup C) \setminus B) \\
 &\quad - f((A \setminus C) \cup B) + f(A \setminus C \setminus B) \leq 0 \quad (4.56)
 \end{aligned}$$

# Submodular Concave

- If the second difference operator everywhere nonpositive:

$$\begin{aligned} f(A \cup B \cup C) - f((A \cup C) \setminus B) \\ - f((A \setminus C) \cup B) + f(A \setminus C \setminus B) \leq 0 \end{aligned} \quad (4.56)$$

then we have the equation:

$$f((A \cup C) \setminus B) + f((A \setminus C) \cup B) \geq f(A \cup B \cup C) + f(A \setminus C \setminus B) \quad (4.57)$$

# Submodular Concave

- If the second difference operator everywhere nonpositive:

$$\begin{aligned} f(A \cup B \cup C) - f((A \cup C) \setminus B) \\ - f((A \setminus C) \cup B) + f(A \setminus C \setminus B) \leq 0 \end{aligned} \quad (4.56)$$

then we have the equation:

$$f((A \cup C) \setminus B) + f((A \setminus C) \cup B) \geq f(A \cup B \cup C) + f(A \setminus C \setminus B) \quad (4.57)$$

- Define  $A' = (A \cup C) \setminus B$  and  $B' = (A \setminus C) \cup B$ . Then the above implies:

$$f(A') + f(B') \geq f(A' \cup B') + f(A' \cap B') \quad (4.58)$$

and note that  $A'$  and  $B'$  so defined can be arbitrary.



# Submodular Concave

- If the second difference operator everywhere nonpositive:

$$f(A \cup B \cup C) - f((A \cup C) \setminus B) - f((A \setminus C) \cup B) + f(A \setminus C \setminus B) \leq 0 \quad (4.56)$$

then we have the equation:

$$f((A \cup C) \setminus B) + f((A \setminus C) \cup B) \geq f(A \cup B \cup C) + f(A \setminus C \setminus B) \quad (4.57)$$


- Define  $A' = (A \cup C) \setminus B$  and  $B' = (A \setminus C) \cup B$ . Then the above implies:

$$f(A') + f(B') \geq f(A' \cup B') + f(A' \cap B') \quad (4.58)$$


and note that  $A'$  and  $B'$  so defined can be arbitrary.

- One sense in which submodular functions are like concave functions.

# Submodular Concave



(a)  $A' = (A \cup C) \setminus B$



(b)  $B' = (A \setminus C) \cup B$

Figure : A figure showing  $A' \cup B' = A \cup B \cup C$  and  $A' \cap B' = A \setminus C \setminus B$ .

# Submodular Concave




Figure : A figure showing  $A' \cup B' = A \cup B \cup C$  and  $A' \cap B' = A \setminus C \setminus B$ .

# Submodularity and Concave

- This submodular/concave relationship is more simply done with singletons.

# Submodularity and Concave

- This submodular/concave relationship is more simply done with singletons.
- Recall four points definition: A function is submodular if for all  $X \subseteq V$  and  $j, k \in V$

$$f(X + j) + f(X + k) \geq f(X + j + k) + f(X) \quad (4.59)$$

# Submodularity and Concave

- This submodular/concave relationship is more simply done with singletons.
- Recall four points definition: A function is submodular if for all  $X \subseteq V$  and  $j, k \in V$

$$f(X + j) + f(X + k) \geq f(X + j + k) + f(X) \quad (4.59)$$

- This gives us a simpler notion corresponding to concavity.

# Submodularity and Concave

- This submodular/concave relationship is more simply done with singletons.
- Recall four points definition: A function is submodular if for all  $X \subseteq V$  and  $j, k \in V$

$$f(X + j) + f(X + k) \geq f(X + j + k) + f(X) \quad (4.59)$$

- This gives us a simpler notion corresponding to concavity.
- Define gain as  $\nabla_j(X) = f(X + j) - f(X)$ , a form of discrete gradient.

# Submodularity and Concave

- This submodular/concave relationship is more simply done with singletons.
- Recall four points definition: A function is submodular if for all  $X \subseteq V$  and  $j, k \in V$

$$f(X + j) + f(X + k) \geq f(X + j + k) + f(X) \quad (4.59)$$

- This gives us a simpler notion corresponding to concavity.
- Define gain as  $\nabla_j(X) = f(X + j) - f(X)$ , a form of discrete gradient.
- Trivially becomes a second-order condition, akin to concave functions: A function is submodular if for all  $X \subseteq V$  and  $j, k \in V$ , we have:

$$\nabla_j \nabla_k f(X) \leq 0 \quad (4.60)$$



# Example: Rank function of a matrix

Consider the following  $4 \times 8$  matrix, so  $V = \{1, 2, 3, 4, 5, 6, 7, 8\}$ .

$$\begin{matrix} & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{pmatrix} 0 & 2 & 2 & 3 & 0 & 1 & 3 & 1 \\ 0 & 3 & 0 & 4 & 0 & 0 & 2 & 4 \\ 0 & 0 & 0 & 0 & 3 & 0 & 0 & 5 \\ 2 & 0 & 0 & 0 & 0 & 0 & 0 & 5 \end{pmatrix} \end{matrix} = \begin{matrix} & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ \begin{pmatrix} | & | & | & | & | & | & | & | \\ x_1 & x_2 & x_3 & x_4 & x_5 & x_6 & x_7 & x_8 \\ | & | & | & | & | & | & | & | \end{pmatrix} \end{matrix}$$

- Let  $A = \{1, 2, 3\}$ ,  $B = \{3, 4, 5\}$ ,  $C = \{6, 7\}$ ,  $A_r = \{1\}$ ,  $B_r = \{5\}$ .
- Then  $r(A) = 3$ ,  $r(B) = 3$ ,  $r(C) = 2$ .
- $r(A \cup C) = 3$ ,  $r(B \cup C) = 3$ .
- $r(A \cup A_r) = 3$ ,  $r(B \cup B_r) = 3$ ,  $r(A \cup B_r) = 4$ ,  $r(B \cup A_r) = 4$ .
- $r(A \cup B) = 4$ ,  $r(A \cap B) = 1 < r(C) = 2$ .
- $6 = r(A) + r(B) > r(A \cup B) + r(A \cap B) = 5$

# On Rank

- Let  $\text{rank} : 2^V \rightarrow \mathbb{Z}_+$  be the rank function.

# On Rank

- Let  $\text{rank} : 2^V \rightarrow \mathbb{Z}_+$  be the rank function.
- In general,  $\text{rank}(A) \leq |A|$ , and vectors in  $A$  are linearly independent if and only if  $\text{rank}(A) = |A|$ .

# On Rank

- Let  $\text{rank} : 2^V \rightarrow \mathbb{Z}_+$  be the rank function.
- In general,  $\text{rank}(A) \leq |A|$ , and vectors in  $A$  are linearly independent if and only if  $\text{rank}(A) = |A|$ .
- If  $A, B$  are such that  $\text{rank}(A) = |A|$  and  $\text{rank}(B) = |B|$ , with  $|A| < |B|$ , then the space spanned by  $B$  is greater, and we can find a vector in  $B$  that is linearly independent of the space spanned by vectors in  $A$ .

# On Rank

- Let  $\text{rank} : 2^V \rightarrow \mathbb{Z}_+$  be the rank function.
- In general,  $\text{rank}(A) \leq |A|$ , and vectors in  $A$  are linearly independent if and only if  $\text{rank}(A) = |A|$ .
- If  $A, B$  are such that  $\text{rank}(A) = |A|$  and  $\text{rank}(B) = |B|$ , with  $|A| < |B|$ , then the space spanned by  $B$  is greater, and we can find a vector in  $B$  that is linearly independent of the space spanned by vectors in  $A$ .
- To stress this point, note that the above condition is  $|A| < |B|$ , **not**  $A \subseteq B$  which is sufficient (to be able to find an independent vector) but not necessary.


# On Rank

- Let  $\text{rank} : 2^V \rightarrow \mathbb{Z}_+$  be the rank function.
- In general,  $\text{rank}(A) \leq |A|$ , and vectors in  $A$  are linearly independent if and only if  $\text{rank}(A) = |A|$ .
- If  $A, B$  are such that  $\text{rank}(A) = |A|$  and  $\text{rank}(B) = |B|$ , with  $|A| < |B|$ , then the space spanned by  $B$  is greater, and we can find a vector in  $B$  that is linearly independent of the space spanned by vectors in  $A$ .
- To stress this point, note that the above condition is  $|A| < |B|$ , **not**  $A \subseteq B$  which is sufficient (to be able to find an independent vector) but not necessary.
- In other words, given  $A, B$  with  $\text{rank}(A) = |A|$  &  $\text{rank}(B) = |B|$ , then  $|A| < |B| \Leftrightarrow \exists \text{ an } b \in B \text{ such that } \text{rank}(A \cup \{b\}) = |A| + 1$ .

# Spanning trees/forests

- We are given a graph  $G = (V, E)$ , and consider the edges  $E = E(G)$  as an index set.
- Consider the  $|V| \times |E|$  incidence matrix of undirected graph  $G$ , which is the matrix  $\mathbf{X}_G = (x_{v,e})_{v \in V(G), e \in E(G)}$  where

$$x_{v,e} = \begin{cases} 1 & \text{if } v \in e \\ 0 & \text{if } v \notin e \end{cases} \quad (4.61)$$



$$\begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 8 \end{matrix} & \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \end{pmatrix} \end{matrix} \quad (4.62)$$

# Spanning trees/forests & incidence matrices

- We are given a graph  $G = (V, E)$ , we can arbitrarily orient the graph (make it directed) consider again the edges  $E = E(G)$  as an index set.
- Consider instead the  $|V| \times |E|$  incidence matrix of undirected graph  $G$ , which is the matrix  $\mathbf{X}_G = (x_{v,e})_{v \in V(G), e \in E(G)}$  where


$$x_{v,e} = \begin{cases} 1 & \text{if } v \in e^+ \\ -1 & \text{if } v \in e^- \\ 0 & \text{if } v \notin e \end{cases} \quad (4.63)$$

and where  $e^+$  is the tail and  $e^-$  is the head of (now) directed edge  $e$ .



# Spanning trees/forests & incidence matrices


- A directed version of the graph (right) and its adjacency matrix (below).
- Orientation can be arbitrary.
- Note, rank of this matrix is 7.



$$\begin{matrix}
 & \begin{matrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 \end{matrix} \\
 \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 8 \end{matrix} & \begin{pmatrix} -1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & -1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 1 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 & 0 & 1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & -1 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 1 & -1 \end{pmatrix}
 \end{matrix}$$

# Spanning trees

- We can consider edge-induced subgraphs and the corresponding matrix columns.




$$\begin{matrix} & 1 \\ 1 & \left( \begin{array}{c} -1 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{array} \right) \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 8 \end{matrix} \quad (4.64)$$

Here,  $\text{rank}(\{x_1\}) = 1$ .

# Spanning trees

- We can consider edge-induced subgraphs and the corresponding matrix columns.




$$\begin{matrix} & \begin{matrix} 1 & 2 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 8 \end{matrix} & \begin{pmatrix} -1 & 1 \\ 1 & 0 \\ 0 & -1 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix} \end{matrix} \quad (4.64)$$

Here,  $\text{rank}(\{x_1, x_2\}) = 2$ .

# Spanning trees

- We can consider edge-induced subgraphs and the corresponding matrix columns.




$$\begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 8 \end{matrix} & \begin{pmatrix} -1 & 1 & 0 \\ 1 & 0 & -1 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \end{matrix} \quad (4.64)$$

Here,  $\text{rank}(\{x_1, x_2, x_3\}) = 3$ .

# Spanning trees

- We can consider edge-induced subgraphs and the corresponding matrix columns.




$$\begin{matrix} & \begin{matrix} 1 & 2 & 3 & 5 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 8 \end{matrix} & \begin{pmatrix} -1 & 1 & 0 & 0 \\ 1 & 0 & -1 & 1 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 \end{pmatrix} \end{matrix} \quad (4.64)$$

Here,  $\text{rank}(\{x_1, x_2, x_3, x_5\}) = 4$ .

# Spanning trees

- We can consider edge-induced subgraphs and the corresponding matrix columns.




$$\begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 & 5 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 8 \end{matrix} & \begin{pmatrix} -1 & 1 & 0 & 0 & 0 \\ 1 & 0 & -1 & 0 & 1 \\ 0 & -1 & 0 & 1 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \end{matrix} \quad (4.64)$$

Here,  $\text{rank}(\{x_1, x_2, x_3, x_4, x_5\}) = 4$ .

# Spanning trees

- We can consider edge-induced subgraphs and the corresponding matrix columns.



$$\begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 8 \end{matrix} & \begin{pmatrix} -1 & 1 & 0 & 0 \\ 1 & 0 & -1 & 0 \\ 0 & -1 & 0 & 1 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \end{matrix} \quad (4.64)$$

Here,  $\text{rank}(\{x_1, x_2, x_3, x_4\}) = 3$  since  $x_4 = -x_1 - x_2 - x_3$ .

# Spanning trees

- In general, whenever the edges specify a cycle, there will be a linear dependence between the corresponding set of vectors in the matrix.
- This means that all forests in the graph correspond to a set of linearly independent column vectors in the matrix.
- Consider a “rank” function defined as follows: given a set of edges  $A \subseteq E(G)$ , the  $\text{rank}(A)$  is the size of the largest forest in the  $A$ -edge induced subgraph of  $G$ .
- The rank of the entire graph then is then a spanning forest of the graph (spanning tree if the graph is connected).
- The rank of the graph is  $\text{rank}(G) = |V| - k$  where  $k$  is the number of connected components of  $G$  (recall, we saw that  $k_G(A)$  is a supermodular function in previous lectures).



# Spanning Tree Algorithms

- We are now given a positive edge-weighted connected graph  $G = (V, E, w)$  where  $w : E \rightarrow \mathbb{R}_+$  is a modular function the edges of the graph. The goal is to find the minimum spanning tree (MST) of the graph.
- Given a tree  $T$ , the cost of the tree is  $\text{cost}(T) = \sum_{e \in T} w(e)$ , the sum of the weights of the edges.
- There are several algorithms for MST:

---

## Algorithm 1: Borůvka's Algorithm

---

```

1  $F \leftarrow \emptyset$  /* We build up the edges of a forest in  $F$  */
2 while  $G(V, F)$  is disconnected do
3   forall the components  $C_i$  of  $F$  do
4      $F \leftarrow F \cup \{e_i\}$  for  $e_i =$  the min-weight edge out of  $C_i$ ;

```

---

# Spanning Tree Algorithms

- We are now given a positive edge-weighted connected graph  $G = (V, E, w)$  where  $w : E \rightarrow \mathbb{R}_+$  is a modular function the edges of the graph. The goal is to find the minimum spanning tree (MST) of the graph.
- Given a tree  $T$ , the cost of the tree is  $\text{cost}(T) = \sum_{e \in T} w(e)$ , the sum of the weights of the edges.
- There are several algorithms for MST:

---

## Algorithm 2: Jarník/Prim/Dijkstra Algorithm

---

- 1  $T \leftarrow \emptyset$  ;
  - 2 **while**  $T$  is not a spanning tree **do**
  - 3      $T \leftarrow T \cup \{e\}$  for  $e =$  the minimum weight edge extending the tree  $T$  to a new vertex ;
-

# Spanning Tree Algorithms

- We are now given a positive edge-weighted connected graph  $G = (V, E, w)$  where  $w : E \rightarrow \mathbb{R}_+$  is a modular function the edges of the graph. The goal is to find the minimum spanning tree (MST) of the graph.
- Given a tree  $T$ , the cost of the tree is  $\text{cost}(T) = \sum_{e \in T} w(e)$ , the sum of the weights of the edges.
- There are several algorithms for MST:

---

## Algorithm 3: Kruskal's Algorithm

---

- 1 Sort the edges so that  $w(e_1) \leq w(e_2) \leq \dots \leq w(e_m)$  ;
  - 2  $T \leftarrow (V(G), \emptyset) = (V, E)$  ;
  - 3 **for**  $i = 1$  **to**  $m$  **do**
  - 4     **if**  $E(T) \cup \{e_i\}$  *does not create a cycle in*  $T$  **then**
  - 5          $E(T) \leftarrow E(T) \cup \{e\}$  ;
-

# Spanning Tree Algorithms

- We are now given a positive edge-weighted connected graph  $G = (V, E, w)$  where  $w : E \rightarrow \mathbb{R}_+$  is a modular function the edges of the graph. The goal is to find the minimum spanning tree (MST) of the graph.
- Given a tree  $T$ , the cost of the tree is  $\text{cost}(T) = \sum_{e \in T} w(e)$ , the sum of the weights of the edges.
- There are several algorithms for MST:
- These three algorithms are all guaranteed to find the optimal minimum spanning tree in (low order) polynomial time.

# Spanning Tree Algorithms

- We are now given a positive edge-weighted connected graph  $G = (V, E, w)$  where  $w : E \rightarrow \mathbb{R}_+$  is a modular function the edges of the graph. The goal is to find the minimum spanning tree (MST) of the graph.
- Given a tree  $T$ , the cost of the tree is  $\text{cost}(T) = \sum_{e \in T} w(e)$ , the sum of the weights of the edges.
- There are several algorithms for MST:
- These three algorithms are all guaranteed to find the optimal minimum spanning tree in (low order) polynomial time.
- These algorithms are **all** related to the “greedy” algorithm. I.e., “add next whatever looks best”.

# Spanning Tree Algorithms

- We are now given a positive edge-weighted connected graph  $G = (V, E, w)$  where  $w : E \rightarrow \mathbb{R}_+$  is a modular function the edges of the graph. The goal is to find the minimum spanning tree (MST) of the graph.
- Given a tree  $T$ , the cost of the tree is  $\text{cost}(T) = \sum_{e \in T} w(e)$ , the sum of the weights of the edges.
- There are several algorithms for MST:
- These three algorithms are all guaranteed to find the optimal minimum spanning tree in (low order) polynomial time.
- These algorithms are **all** related to the “greedy” algorithm. I.e., “add next whatever looks best”.
- These algorithms will also always find a basis (a set of linearly independent vectors that span the underlying space) in the matrix example we saw earlier.

# Spanning Tree Algorithms

- We are now given a positive edge-weighted connected graph  $G = (V, E, w)$  where  $w : E \rightarrow \mathbb{R}_+$  is a modular function the edges of the graph. The goal is to find the minimum spanning tree (MST) of the graph.
- Given a tree  $T$ , the cost of the tree is  $\text{cost}(T) = \sum_{e \in T} w(e)$ , the sum of the weights of the edges.
- There are several algorithms for MST:
- These three algorithms are all guaranteed to find the optimal minimum spanning tree in (low order) polynomial time.
- These algorithms are **all** related to the “greedy” algorithm. I.e., “add next whatever looks best”.
- These algorithms will also always find a basis (a set of linearly independent vectors that span the underlying space) in the matrix example we saw earlier.
- The above are all examples of a matroid, which is the fundamental reason why the greedy algorithms work.

# From Matrix Rank $\rightarrow$ Matroid

- So  $V$  is set of column vector indices of a matrix.



# From Matrix Rank $\rightarrow$ Matroid

- So  $V$  is set of column vector indices of a matrix.
- Let  $\mathcal{I}$  be a set of all subsets of  $V$  such that for any  $I \in \mathcal{I}$ , the vectors indexed by  $I$  are linearly independent.

# From Matrix Rank $\rightarrow$ Matroid

- So  $V$  is set of column vector indices of a matrix.
- Let  $\mathcal{I}$  be a set of all subsets of  $V$  such that for any  $I \in \mathcal{I}$ , the vectors indexed by  $I$  are linearly independent.
- Given a set  $B \in \mathcal{I}$  of linearly independent vectors, then any subset  $A \subseteq B$  is also linearly independent.

# From Matrix Rank $\rightarrow$ Matroid

- So  $V$  is set of column vector indices of a matrix.
- Let  $\mathcal{I}$  be a set of all subsets of  $V$  such that for any  $I \in \mathcal{I}$ , the vectors indexed by  $I$  are linearly independent.
- Given a set  $B \in \mathcal{I}$  of linearly independent vectors, then any subset  $A \subseteq B$  is also linearly independent. Hence,  $\mathcal{I}$  is down-closed or “subclusive”, under subsets.

# From Matrix Rank $\rightarrow$ Matroid

- So  $V$  is set of column vector indices of a matrix.
- Let  $\mathcal{I}$  be a set of all subsets of  $V$  such that for any  $I \in \mathcal{I}$ , the vectors indexed by  $I$  are linearly independent.
- Given a set  $B \in \mathcal{I}$  of linearly independent vectors, then any subset  $A \subseteq B$  is also linearly independent. Hence,  $\mathcal{I}$  is down-closed or “**subclusive**”, under subsets. In other words,

$$A \subseteq B \text{ and } B \in \mathcal{I} \Rightarrow A \in \mathcal{I} \quad (4.65)$$

# From Matrix Rank $\rightarrow$ Matroid

- So  $V$  is set of column vector indices of a matrix.
- Let  $\mathcal{I}$  be a set of all subsets of  $V$  such that for any  $I \in \mathcal{I}$ , the vectors indexed by  $I$  are linearly independent.
- Given a set  $B \in \mathcal{I}$  of linearly independent vectors, then any subset  $A \subseteq B$  is also linearly independent. Hence,  $\mathcal{I}$  is down-closed or “**subclusive**”, under subsets. In other words,

$$A \subseteq B \text{ and } B \in \mathcal{I} \Rightarrow A \in \mathcal{I} \quad (4.65)$$

- **maxInd**: Inclusionwise maximal independent subsets (or **bases**) of  $B$ .

$$\text{maxInd}(B) \triangleq \{A \subseteq B : A \in \mathcal{I} \text{ and } \forall v \in B \setminus A, A \cup \{v\} \notin \mathcal{I}\} \quad (4.66)$$

# From Matrix Rank $\rightarrow$ Matroid

- So  $V$  is set of column vector indices of a matrix.
- Let  $\mathcal{I}$  be a set of all subsets of  $V$  such that for any  $I \in \mathcal{I}$ , the vectors indexed by  $I$  are linearly independent.
- Given a set  $B \in \mathcal{I}$  of linearly independent vectors, then any subset  $A \subseteq B$  is also linearly independent. Hence,  $\mathcal{I}$  is down-closed or “**subclusive**”, under subsets. In other words,

$$A \subseteq B \text{ and } B \in \mathcal{I} \Rightarrow A \in \mathcal{I} \quad (4.65)$$

- **maxInd**: Inclusionwise maximal independent subsets (or **bases**) of  $B$ .

$$\text{maxInd}(B) \triangleq \{A \subseteq B : A \in \mathcal{I} \text{ and } \forall v \in B \setminus A, A \cup \{v\} \notin \mathcal{I}\} \quad (4.66)$$

- Given any set  $B \subset V$  of vectors, all maximal (by set inclusion) subsets of linearly independent vectors are the same size. That is, for all  $B \subseteq V$ ,

$$\forall A_1, A_2 \in \text{maxInd}(B), \quad |A_1| = |A_2| \quad (4.67)$$

# From Matrix Rank $\rightarrow$ Matroid

- Thus, for all  $I \in \mathcal{I}$ , the matrix rank function has the property

$$r(I) = |I| \quad (4.68)$$

and for any  $B \notin \mathcal{I}$ ,

$$r(B) = \max \{|A| : A \subseteq B \text{ and } A \in \mathcal{I}\} \leq |B| \quad (4.69)$$

# Matroid

- Matroids abstract the notion of linear independence of a set of vectors to general algebraic properties.



# Matroid

- Matroids abstract the notion of linear independence of a set of vectors to general algebraic properties.
- In a matroid, there is an underlying **ground set**, say  $E$  (or  $V$ ), and a collection of subsets of  $E$  that correspond to independent elements.

# Matroid

- Matroids abstract the notion of linear independence of a set of vectors to general algebraic properties.
- In a matroid, there is an underlying **ground set**, say  $E$  (or  $V$ ), and a collection of subsets of  $E$  that correspond to independent elements.
- There are many definitions of matroids that are mathematically equivalent, we'll see some of them here.

# Independence System

## Definition 4.5.1 (set system)

A (finite) ground set  $E$  and a set of subsets of  $E$ ,  $\emptyset \neq \mathcal{I} \subseteq 2^E$  is called a set system, notated  $(E, \mathcal{I})$ .

- Set systems can be arbitrarily complex since, as stated, there is no method to determine if a given set  $S \subseteq E$  has  $S \in \mathcal{I}$ .

# Independence System

## Definition 4.5.1 (set system)

A (finite) ground set  $E$  and a set of subsets of  $E$ ,  $\emptyset \neq \mathcal{I} \subseteq 2^E$  is called a set system, notated  $(E, \mathcal{I})$ .

- Set systems can be arbitrarily complex since, as stated, there is no method to determine if a given set  $S \subseteq E$  has  $S \in \mathcal{I}$ .
- One useful property is “heredity.” Namely, a set system is a hereditary set system if for any  $A \subset B \in \mathcal{I}$ , we have that  $A \in \mathcal{I}$ .

# Independence System

## Definition 4.5.2 (independence (or hereditary) system)

A set system  $(V, \mathcal{I})$  is an independence system if

$$\emptyset \in \mathcal{I} \quad (\text{emptyset containing}) \quad (I1)$$

and

$$\forall I \in \mathcal{I}, J \subset I \Rightarrow J \in \mathcal{I} \quad (\text{subclusive}) \quad (I2)$$

- Property I2 is called “down monotone,” “down closed,” or “subclusive”

# Independence System

## Definition 4.5.2 (independence (or hereditary) system)

A set system  $(V, \mathcal{I})$  is an independence system if

$$\emptyset \in \mathcal{I} \quad (\text{emptyset containing}) \quad (I1)$$

and

$$\forall I \in \mathcal{I}, J \subset I \Rightarrow J \in \mathcal{I} \quad (\text{subclusive}) \quad (I2)$$

- Property I2 is called “down monotone,” “down closed,” or “subclusive”
- Example:  $E = \{1, 2, 3, 4\}$ . With  $\mathcal{I} = \{\emptyset, \{1\}, \{1, 2\}, \{1, 2, 4\}\}$ .

# Independence System

## Definition 4.5.2 (independence (or hereditary) system)

A set system  $(V, \mathcal{I})$  is an independence system if

$$\emptyset \in \mathcal{I} \quad (\text{emptyset containing}) \quad (I1)$$

and

$$\forall I \in \mathcal{I}, J \subset I \Rightarrow J \in \mathcal{I} \quad (\text{subclusive}) \quad (I2)$$

- Property I2 is called “down monotone,” “down closed,” or “subclusive”
- Example:  $E = \{1, 2, 3, 4\}$ . With  $\mathcal{I} = \{\emptyset, \{1\}, \{1, 2\}, \{1, 2, 4\}\}$ .
- Then  $(E, \mathcal{I})$  is a set system, but not an independence system since it is not down closed (i.e., we have  $\{1, 2\} \in \mathcal{I}$  but not  $\{2\} \in \mathcal{I}$ ).

# Independence System

## Definition 4.5.2 (independence (or hereditary) system)

A set system  $(V, \mathcal{I})$  is an independence system if

$$\emptyset \in \mathcal{I} \quad (\text{emptyset containing}) \quad (I1)$$

and

$$\forall I \in \mathcal{I}, J \subset I \Rightarrow J \in \mathcal{I} \quad (\text{subclusive}) \quad (I2)$$

- Property I2 is called “down monotone,” “down closed,” or “subclusive”
- Example:  $E = \{1, 2, 3, 4\}$ . With  $\mathcal{I} = \{\emptyset, \{1\}, \{1, 2\}, \{1, 2, 4\}\}$ .
- Then  $(E, \mathcal{I})$  is a set system, but not an independence system since it is not down closed (i.e., we have  $\{1, 2\} \in \mathcal{I}$  but not  $\{2\} \in \mathcal{I}$ ).
- With  $\mathcal{I} = \{\emptyset, \{1\}, \{2\}, \{1, 2\}\}$ , then  $(E, \mathcal{I})$  is now an independence (hereditary) system.



# Independence System

$$\begin{array}{c}
 \begin{array}{cccccccc}
 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\
 1 & \left( \begin{array}{cccccccc}
 0 & 0 & 1 & 1 & 2 & 1 & 3 & 1
 \end{array} \right) \\
 2 & \left( \begin{array}{cccccccc}
 0 & 1 & 1 & 0 & 2 & 0 & 2 & 4
 \end{array} \right) \\
 3 & \left( \begin{array}{cccccccc}
 1 & 1 & 1 & 0 & 0 & 3 & 1 & 5
 \end{array} \right)
 \end{array}
 =
 \begin{array}{c}
 \begin{array}{cccccccc}
 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\
 \left( \begin{array}{cccccccc}
 | & | & | & | & | & | & | & | \\
 x_1 & x_2 & x_3 & x_4 & x_5 & x_6 & x_7 & x_8 \\
 | & | & | & | & | & | & | & |
 \end{array} \right)
 \end{array}
 \end{array}
 \quad (4.70)$$

- Given any set of linearly independent vectors  $A$ , any subset  $B \subset A$  will also be linearly independent.

# Independence System

$$\begin{array}{c}
 \begin{array}{cccccccc}
 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\
 1 & \left( \begin{array}{cccccccc}
 0 & 0 & 1 & 1 & 2 & 1 & 3 & 1
 \end{array} \right) \\
 2 & \left( \begin{array}{cccccccc}
 0 & 1 & 1 & 0 & 2 & 0 & 2 & 4
 \end{array} \right) \\
 3 & \left( \begin{array}{cccccccc}
 1 & 1 & 1 & 0 & 0 & 3 & 1 & 5
 \end{array} \right)
 \end{array}
 =
 \begin{array}{c}
 \begin{array}{cccccccc}
 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\
 \left( \begin{array}{cccccccc}
 | & | & | & | & | & | & | & | \\
 x_1 & x_2 & x_3 & x_4 & x_5 & x_6 & x_7 & x_8 \\
 | & | & | & | & | & | & | & |
 \end{array} \right)
 \end{array}
 \end{array}
 \quad (4.70)$$

- Given any set of linearly independent vectors  $A$ , any subset  $B \subset A$  will also be linearly independent.
- Given any forest  $G_f$  that is an edge-induced sub-graph of a graph  $G$ , any sub-graph of  $G_f$  is also a forest.

# Independence System

$$\begin{matrix} & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ \begin{matrix} 1 \\ 2 \\ 3 \end{matrix} & \begin{pmatrix} 0 & 0 & 1 & 1 & 2 & 1 & 3 & 1 \\ 0 & 1 & 1 & 0 & 2 & 0 & 2 & 4 \\ 1 & 1 & 1 & 0 & 0 & 3 & 1 & 5 \end{pmatrix} \end{matrix} = \begin{matrix} & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ \begin{pmatrix} | & | & | & | & | & | & | & | \\ x_1 & x_2 & x_3 & x_4 & x_5 & x_6 & x_7 & x_8 \\ | & | & | & | & | & | & | & | \end{pmatrix} \end{matrix} \quad (4.70)$$

- Given any set of linearly independent vectors  $A$ , any subset  $B \subset A$  will also be linearly independent.
- Given any forest  $G_f$  that is an edge-induced sub-graph of a graph  $G$ , any sub-graph of  $G_f$  is also a forest.
- So these both constitute independence systems.

# Matroid

Independent set definition of a matroid is perhaps most natural. Note, if  $J \in \mathcal{I}$ , then  $J$  is said to be an **independent set**.

## Definition 4.5.3 (Matroid)

A set system  $(E, \mathcal{I})$  is a **Matroid** if

- (I1)  $\emptyset \in \mathcal{I}$
- (I2)  $\forall I \in \mathcal{I}, J \subset I \Rightarrow J \in \mathcal{I}$
- (I3)  $\forall I, J \in \mathcal{I}$ , with  $|I| = |J| + 1$ , then there exists  $x \in I \setminus J$  such that  $J \cup \{x\} \in \mathcal{I}$ .

# Matroid

Slight modification (non unit increment) that is equivalent.

## Definition 4.5.4 (Matroid-II)

A set system  $(E, \mathcal{I})$  is a **Matroid** if

- (I1')  $\emptyset \in \mathcal{I}$
- (I2')  $\forall I \in \mathcal{I}, J \subset I \Rightarrow J \in \mathcal{I}$  (or “down-closed”)
- (I3')  $\forall I, J \in \mathcal{I}$ , with  $|I| > |J|$ , then there exists  $x \in I \setminus J$  such that  $J \cup \{x\} \in \mathcal{I}$

Note (I1)=(I1'), (I2)=(I2'), and we get (I3)≡(I3') using induction.

# Matroids, independent sets, and bases

- **Independent sets:** Given a matroid  $M = (E, \mathcal{I})$ , a subset  $A \subseteq E$  is called **independent** if  $A \in \mathcal{I}$  and otherwise  $A$  is called **dependent**.

# Matroids, independent sets, and bases

- **Independent sets:** Given a matroid  $M = (E, \mathcal{I})$ , a subset  $A \subseteq E$  is called **independent** if  $A \in \mathcal{I}$  and otherwise  $A$  is called **dependent**.
- **A base of  $U \subseteq E$ :** For  $U \subseteq E$ , a subset  $B \subseteq U$  is called a **base** of  $U$  if  $B$  is inclusionwise maximally independent subset of  $U$ . That is,  $B \in \mathcal{I}$  and there is no  $Z \in \mathcal{I}$  with  $B \subset Z \subseteq U$ .

# Matroids, independent sets, and bases

- **Independent sets:** Given a matroid  $M = (E, \mathcal{I})$ , a subset  $A \subseteq E$  is called **independent** if  $A \in \mathcal{I}$  and otherwise  $A$  is called **dependent**.
- **A base of  $U \subseteq E$ :** For  $U \subseteq E$ , a subset  $B \subseteq U$  is called a **base** of  $U$  if  $B$  is inclusionwise maximally independent subset of  $U$ . That is,  $B \in \mathcal{I}$  and there is no  $Z \in \mathcal{I}$  with  $B \subset Z \subseteq U$ .
- **A base of a matroid:** If  $U = E$ , then a “base of  $E$ ” is just called a **base** of the matroid  $M$  (this corresponds to a **basis** in a linear space).



# Matroids - important property

## Proposition 4.5.5

*In a matroid  $M = (E, \mathcal{I})$ , for any  $U \subseteq E(M)$ , any two bases of  $U$  have the same size.*

# Matroids - important property

## Proposition 4.5.5

*In a matroid  $M = (E, \mathcal{I})$ , for any  $U \subseteq E(M)$ , any two bases of  $U$  have the same size.*

- In matrix terms, given a set of vectors  $U$ , all sets of independent vectors that span the space spanned by  $U$  have the same size.

# Matroids - important property

## Proposition 4.5.5

*In a matroid  $M = (E, \mathcal{I})$ , for any  $U \subseteq E(M)$ , any two bases of  $U$  have the same size.*

- In matrix terms, given a set of vectors  $U$ , all sets of independent vectors that span the space spanned by  $U$  have the same size.
- In fact, under (I1),(I2), this condition is equivalent to (I3). **Exercise:** show the following is equivalent to the above.

# Matroids - important property

## Proposition 4.5.5

*In a matroid  $M = (E, \mathcal{I})$ , for any  $U \subseteq E(M)$ , any two bases of  $U$  have the same size.*

- In matrix terms, given a set of vectors  $U$ , all sets of independent vectors that span the space spanned by  $U$  have the same size.
- In fact, under (I1), (I2), this condition is equivalent to (I3). **Exercise:** show the following is equivalent to the above.

## Definition 4.5.6 (Matroid)

A set system  $(V, \mathcal{I})$  is a **Matroid** if

# Matroids - important property

## Proposition 4.5.5

*In a matroid  $M = (E, \mathcal{I})$ , for any  $U \subseteq E(M)$ , any two bases of  $U$  have the same size.*

- In matrix terms, given a set of vectors  $U$ , all sets of independent vectors that span the space spanned by  $U$  have the same size.
- In fact, under (I1), (I2), this condition is equivalent to (I3). **Exercise:** show the following is equivalent to the above.

## Definition 4.5.6 (Matroid)

A set system  $(V, \mathcal{I})$  is a **Matroid** if

(I1')  $\emptyset \in \mathcal{I}$  (emptyset containing)

# Matroids - important property

## Proposition 4.5.5

*In a matroid  $M = (E, \mathcal{I})$ , for any  $U \subseteq E(M)$ , any two bases of  $U$  have the same size.*

- In matrix terms, given a set of vectors  $U$ , all sets of independent vectors that span the space spanned by  $U$  have the same size.
- In fact, under (I1), (I2), this condition is equivalent to (I3). **Exercise:** show the following is equivalent to the above.

## Definition 4.5.6 (Matroid)

A set system  $(V, \mathcal{I})$  is a **Matroid** if

(I1')  $\emptyset \in \mathcal{I}$  (emptyset containing)

(I2')  $\forall I \in \mathcal{I}, J \subset I \Rightarrow J \in \mathcal{I}$  (down-closed or subclusive)

# Matroids - important property

## Proposition 4.5.5

*In a matroid  $M = (E, \mathcal{I})$ , for any  $U \subseteq E(M)$ , any two bases of  $U$  have the same size.*

- In matrix terms, given a set of vectors  $U$ , all sets of independent vectors that span the space spanned by  $U$  have the same size.
- In fact, under (I1), (I2), this condition is equivalent to (I3). **Exercise:** show the following is equivalent to the above.

## Definition 4.5.6 (Matroid)

A set system  $(V, \mathcal{I})$  is a **Matroid** if

- (I1')  $\emptyset \in \mathcal{I}$  (emptyset containing)
- (I2')  $\forall I \in \mathcal{I}, J \subset I \Rightarrow J \in \mathcal{I}$  (down-closed or subclusive)
- (I3')  $\forall X \subseteq V$ , and  $I_1, I_2 \in \max \text{Ind}(X)$ , we have  $|I_1| = |I_2|$  (all maximally independent subsets of  $X$  have the same size).

# Matroids - rank

- Recall, in any matroid  $M = (E, \mathcal{I})$ ,  $\forall U \subseteq E(M)$ , any two bases of  $U$  have the same size.



# Matroids - rank

- Recall, in any matroid  $M = (E, \mathcal{I})$ ,  $\forall U \subseteq E(M)$ , any two bases of  $U$  have the same size.
- The common size of all the **bases** of  $U$  is called the rank of  $U$ , denoted  $r_M(U)$  or just  $r(U)$  when the matroid in equation is unambiguous.

# Matroids - rank

- Recall, in any matroid  $M = (E, \mathcal{I})$ ,  $\forall U \subseteq E(M)$ , any two bases of  $U$  have the same size.
- The common size of all the **bases** of  $U$  is called the rank of  $U$ , denoted  $r_M(U)$  or just  $r(U)$  when the matroid in equation is unambiguous.
- $r(E) = r_{(E, \mathcal{I})}$  is the rank of the matroid, and is the common size of all the bases of the matroid.

# Matroids - rank

- Recall, in any matroid  $M = (E, \mathcal{I})$ ,  $\forall U \subseteq E(M)$ , any two bases of  $U$  have the same size.
- The common size of all the **bases** of  $U$  is called the rank of  $U$ , denoted  $r_M(U)$  or just  $r(U)$  when the matroid in equation is unambiguous.
- $r(E) = r_{(E, \mathcal{I})}$  is the rank of the matroid, and is the common size of all the bases of the matroid.
- We can a bit more formally define the rank function this way.

# Matroids - rank

- Recall, in any matroid  $M = (E, \mathcal{I})$ ,  $\forall U \subseteq E(M)$ , any two bases of  $U$  have the same size.
- The common size of all the **bases** of  $U$  is called the rank of  $U$ , denoted  $r_M(U)$  or just  $r(U)$  when the matroid in equation is unambiguous.
- $r(E) = r_{(E, \mathcal{I})}$  is the rank of the matroid, and is the common size of all the bases of the matroid.
- We can a bit more formally define the rank function this way.

## Definition 4.5.7 (matroid rank function)

The rank of a matroid is a function  $r : 2^E \rightarrow \mathbb{Z}_+$  defined by

$$r(A) = \max \{|X| : X \subseteq A, X \in \mathcal{I}\} = \max_{X \in \mathcal{I}} |A \cap X| \quad (4.71)$$

# Matroids - rank

- Recall, in any matroid  $M = (E, \mathcal{I})$ ,  $\forall U \subseteq E(M)$ , any two bases of  $U$  have the same size.
- The common size of all the **bases** of  $U$  is called the rank of  $U$ , denoted  $r_M(U)$  or just  $r(U)$  when the matroid in equation is unambiguous.
- $r(E) = r_{(E, \mathcal{I})}$  is the rank of the matroid, and is the common size of all the bases of the matroid.
- We can a bit more formally define the rank function this way.

## Definition 4.5.7 (matroid rank function)

The rank of a matroid is a function  $r : 2^E \rightarrow \mathbb{Z}_+$  defined by

$$r(A) = \max \{|X| : X \subseteq A, X \in \mathcal{I}\} = \max_{X \in \mathcal{I}} |A \cap X| \quad (4.71)$$

- From the above, we immediately see that  $r(A) \leq |A|$ .

# Matroids - rank

- Recall, in any matroid  $M = (E, \mathcal{I})$ ,  $\forall U \subseteq E(M)$ , any two bases of  $U$  have the same size.
- The common size of all the **bases** of  $U$  is called the rank of  $U$ , denoted  $r_M(U)$  or just  $r(U)$  when the matroid in equation is unambiguous.
- $r(E) = r_{(E, \mathcal{I})}$  is the rank of the matroid, and is the common size of all the bases of the matroid.
- We can a bit more formally define the rank function this way.

## Definition 4.5.7 (matroid rank function)

The rank of a matroid is a function  $r : 2^E \rightarrow \mathbb{Z}_+$  defined by

$$r(A) = \max \{|X| : X \subseteq A, X \in \mathcal{I}\} = \max_{X \in \mathcal{I}} |A \cap X| \quad (4.71)$$

- From the above, we immediately see that  $r(A) \leq |A|$ .
- Moreover, if  $r(A) = |A|$ , then  $A \in \mathcal{I}$ , meaning  $A$  is independent (in this case,  $A$  is a **self base**).

# Matroids, other definitions using matroid rank $r : 2^V \rightarrow \mathbb{Z}_+$

## Definition 4.5.8 (closed/flat/subspace)

A subset  $A \subseteq E$  is **closed** (equivalently, a **flat** or a **subspace**) of matroid  $M$  if for all  $x \in E \setminus A$ ,  $r(A \cup \{x\}) = r(A) + 1$ .

# Matroids, other definitions using matroid rank $r : 2^V \rightarrow \mathbb{Z}_+$

## Definition 4.5.8 (closed/flat/subspace)

A subset  $A \subseteq E$  is **closed** (equivalently, a **flat** or a **subspace**) of matroid  $M$  if for all  $x \in E \setminus A$ ,  $r(A \cup \{x\}) = r(A) + 1$ .

## Definition 4.5.9 (closure)

Given  $A \subseteq E$ , the **closure** (or **span**) of  $A$ , is defined by  $\text{span}(A) = \{b \in E : r(A \cup \{b\}) = r(A)\}$ .



# Matroids, other definitions using matroid rank $r : 2^V \rightarrow \mathbb{Z}_+$

## Definition 4.5.8 (closed/flat/subspace)

A subset  $A \subseteq E$  is **closed** (equivalently, a **flat** or a **subspace**) of matroid  $M$  if for all  $x \in E \setminus A$ ,  $r(A \cup \{x\}) = r(A) + 1$ .

## Definition 4.5.9 (closure)

Given  $A \subseteq E$ , the **closure** (or **span**) of  $A$ , is defined by  $\text{span}(A) = \{b \in E : r(A \cup \{b\}) = r(A)\}$ .

Therefore, a closed set  $A$  has  $\text{span}(A) = A$ .

# Matroids, other definitions using matroid rank $r : 2^V \rightarrow \mathbb{Z}_+$

## Definition 4.5.8 (closed/flat/subspace)

A subset  $A \subseteq E$  is **closed** (equivalently, a **flat** or a **subspace**) of matroid  $M$  if for all  $x \in E \setminus A$ ,  $r(A \cup \{x\}) = r(A) + 1$ .

## Definition 4.5.9 (closure)

Given  $A \subseteq E$ , the **closure** (or **span**) of  $A$ , is defined by  $\text{span}(A) = \{b \in E : r(A \cup \{b\}) = r(A)\}$ .

Therefore, a closed set  $A$  has  $\text{span}(A) = A$ .

## Definition 4.5.10 (circuit)

A subset  $A \subseteq E$  is **circuit** or a **cycle** if it is an inclusionwise-minimal dependent set (i.e., if  $r(A) < |A|$  and for any  $a \in A$ ,  $r(A \setminus \{a\}) = |A| - 1$ ).

# Matroids by bases

In general, besides independent sets and rank functions, there are other equivalent ways to characterize matroids.

## Theorem 4.5.11 (Matroid (by bases))

*Let  $E$  be a set and  $\mathcal{B}$  be a nonempty collection of subsets of  $E$ . Then the following are equivalent.*

- ①  *$\mathcal{B}$  is the collection of bases of a matroid;*
- ② *if  $B, B' \in \mathcal{B}$ , and  $x \in B' \setminus B$ , then  $B' - x + y \in \mathcal{B}$  for some  $y \in B \setminus B'$ .*
- ③ *If  $B, B' \in \mathcal{B}$ , and  $x \in B' \setminus B$ , then  $B - y + x \in \mathcal{B}$  for some  $y \in B \setminus B'$ .*

Properties 2 and 3 are called “exchange properties.”

# Matroids by bases

In general, besides independent sets and rank functions, there are other equivalent ways to characterize matroids.

## Theorem 4.5.11 (Matroid (by bases))

*Let  $E$  be a set and  $\mathcal{B}$  be a nonempty collection of subsets of  $E$ . Then the following are equivalent.*

- ①  $\mathcal{B}$  is the collection of bases of a matroid;
- ② if  $B, B' \in \mathcal{B}$ , and  $x \in B' \setminus B$ , then  $B' - x + y \in \mathcal{B}$  for some  $y \in B \setminus B'$ .
- ③ If  $B, B' \in \mathcal{B}$ , and  $x \in B' \setminus B$ , then  $B - y + x \in \mathcal{B}$  for some  $y \in B \setminus B'$ .

Properties 2 and 3 are called “exchange properties.”

Proof here is omitted but think about this for a moment in terms of linear spaces and matrices, and (alternatively) spanning trees.

# Matroids by circuits

A set is independent if and only if it contains no circuit. Therefore, it is not surprising that circuits can also characterize a matroid.

## Theorem 4.5.12 (Matroid by circuits)

*Let  $E$  be a set and  $\mathcal{C}$  be a collection of subsets of  $E$  that satisfy the following three properties:*

- ① (C1):  $\emptyset \notin \mathcal{C}$
- ② (C2): if  $C_1, C_2 \in \mathcal{C}$  and  $C_1 \subseteq C_2$ , then  $C_1 = C_2$ .
- ③ (C3): if  $C_1, C_2 \in \mathcal{C}$  with  $C_1 \neq C_2$ , and  $C \in C_1 \cap C_2$ , then there exists a  $C_3 \in \mathcal{C}$  such that  $C_3 \subseteq (C_1 \cup C_2) \setminus \{e\}$ .

# Matroids by circuits

Several circuit definitions for matroids.

## Theorem 4.5.13 (Matroid by circuits)

*Let  $E$  be a set and  $\mathcal{C}$  be a collection of nonempty subsets of  $E$ , such that no two sets in  $\mathcal{C}$  are contained in each other. Then the following are equivalent.*

- ①  *$\mathcal{C}$  is the collection of circuits of a matroid;*
- ② *if  $C, C' \in \mathcal{C}$ , and  $x \in C \cap C'$ , then  $(C \cup C') \setminus \{x\}$  contains a set in  $\mathcal{C}$ ;*
- ③ *if  $C, C' \in \mathcal{C}$ , and  $x \in C \cap C'$ , and  $y \in C \setminus C'$ , then  $(C \cup C') \setminus \{x\}$  contains a set in  $\mathcal{C}$  containing  $y$ ;*

# Matroids by circuits

Several circuit definitions for matroids.

## Theorem 4.5.13 (Matroid by circuits)

*Let  $E$  be a set and  $\mathcal{C}$  be a collection of nonempty subsets of  $E$ , such that no two sets in  $\mathcal{C}$  are contained in each other. Then the following are equivalent.*

- ①  *$\mathcal{C}$  is the collection of circuits of a matroid;*
- ② *if  $C, C' \in \mathcal{C}$ , and  $x \in C \cap C'$ , then  $(C \cup C') \setminus \{x\}$  contains a set in  $\mathcal{C}$ ;*
- ③ *if  $C, C' \in \mathcal{C}$ , and  $x \in C \cap C'$ , and  $y \in C \setminus C'$ , then  $(C \cup C') \setminus \{x\}$  contains a set in  $\mathcal{C}$  containing  $y$ ;*

Again, think about this for a moment in terms of linear spaces and matrices, and spanning trees.

# Matroids by submodular functions

## Theorem 4.5.14 (Matroid by submodular functions)

Let  $f : 2^E \rightarrow \mathbb{Z}$  be a integer valued monotone non-decreasing submodular function. Define a set of sets as follows:

$$\mathcal{C}(f) = \left\{ C \subseteq E : \begin{array}{l} C \text{ is non-empty,} \\ \text{is inclusionwise-minimal,} \\ \text{and has } f(C) < |C| \end{array} \right\} \quad (4.72)$$

Then  $\mathcal{C}(f)$  is the collection of circuits of a matroid on  $E$ .

Inclusionwise-minimal means that if  $C \in \mathcal{C}(f)$ , then there exists no  $C' \subset C$  with  $C' \in \mathcal{C}(f)$  (i.e.,  $C' \subset C$  would either be empty or have  $|C'| \leq f(C')$ ).



# Uniform Matroid

- Given  $E$ , consider  $\mathcal{I}$  to be all subsets of  $E$  that are at most size  $k$ .  
That is  $\mathcal{I} = \{A \subseteq E : |A| \leq k\}$ .

# Uniform Matroid

- Given  $E$ , consider  $\mathcal{I}$  to be all subsets of  $E$  that are at most size  $k$ .  
That is  $\mathcal{I} = \{A \subseteq E : |A| \leq k\}$ .
- Then  $(E, \mathcal{I})$  is a matroid called a  $k$ -uniform matroid.

# Uniform Matroid

- Given  $E$ , consider  $\mathcal{I}$  to be all subsets of  $E$  that are at most size  $k$ .  
That is  $\mathcal{I} = \{A \subseteq E : |A| \leq k\}$ .
- Then  $(E, \mathcal{I})$  is a matroid called a  $k$ -uniform matroid.
- Note, if  $I, J \in \mathcal{I}$ , and  $|I| < |J| \leq k$ , and  $j \in J$  such that  $j \notin I$ , then  $j$  is such that  $|I + j| \leq k$  and so  $I + j \in \mathcal{I}$ .

# Uniform Matroid

- Given  $E$ , consider  $\mathcal{I}$  to be all subsets of  $E$  that are at most size  $k$ .  
That is  $\mathcal{I} = \{A \subseteq E : |A| \leq k\}$ .
- Then  $(E, \mathcal{I})$  is a matroid called a  $k$ -uniform matroid.
- Note, if  $I, J \in \mathcal{I}$ , and  $|I| < |J| \leq k$ , and  $j \in J$  such that  $j \notin I$ , then  $j$  is such that  $|I + j| \leq k$  and so  $I + j \in \mathcal{I}$ .
- Rank function

$$r(A) = \begin{cases} |A| & \text{if } |A| \leq k \\ k & \text{if } |A| > k \end{cases} \quad (4.73)$$

# Uniform Matroid

- Given  $E$ , consider  $\mathcal{I}$  to be all subsets of  $E$  that are at most size  $k$ . That is  $\mathcal{I} = \{A \subseteq E : |A| \leq k\}$ .
- Then  $(E, \mathcal{I})$  is a matroid called a  $k$ -uniform matroid.
- Note, if  $I, J \in \mathcal{I}$ , and  $|I| < |J| \leq k$ , and  $j \in J$  such that  $j \notin I$ , then  $j$  is such that  $|I + j| \leq k$  and so  $I + j \in \mathcal{I}$ .
- Rank function

$$r(A) = \begin{cases} |A| & \text{if } |A| \leq k \\ k & \text{if } |A| > k \end{cases} \quad (4.73)$$

- Note, this function is submodular. Not surprising since  $r(A) = \min(|A|, k)$  which is a non-decreasing concave function applied to a modular function.

# Uniform Matroid

- Given  $E$ , consider  $\mathcal{I}$  to be all subsets of  $E$  that are at most size  $k$ . That is  $\mathcal{I} = \{A \subseteq E : |A| \leq k\}$ .
- Then  $(E, \mathcal{I})$  is a matroid called a  $k$ -uniform matroid.
- Note, if  $I, J \in \mathcal{I}$ , and  $|I| < |J| \leq k$ , and  $j \in J$  such that  $j \notin I$ , then  $j$  is such that  $|I + j| \leq k$  and so  $I + j \in \mathcal{I}$ .
- Rank function

$$r(A) = \begin{cases} |A| & \text{if } |A| \leq k \\ k & \text{if } |A| > k \end{cases} \quad (4.73)$$

- Note, this function is submodular. Not surprising since  $r(A) = \min(|A|, k)$  which is a non-decreasing concave function applied to a modular function.
- Closure function

$$\text{span}(A) = \begin{cases} A & \text{if } |A| < k, \\ E & \text{if } |A| \geq k, \end{cases} \quad (4.74)$$

# Uniform Matroid

- Given  $E$ , consider  $\mathcal{I}$  to be all subsets of  $E$  that are at most size  $k$ . That is  $\mathcal{I} = \{A \subseteq E : |A| \leq k\}$ .
- Then  $(E, \mathcal{I})$  is a matroid called a  $k$ -uniform matroid.
- Note, if  $I, J \in \mathcal{I}$ , and  $|I| < |J| \leq k$ , and  $j \in J$  such that  $j \notin I$ , then  $j$  is such that  $|I + j| \leq k$  and so  $I + j \in \mathcal{I}$ .
- Rank function

$$r(A) = \begin{cases} |A| & \text{if } |A| \leq k \\ k & \text{if } |A| > k \end{cases} \quad (4.73)$$

- Note, this function is submodular. Not surprising since  $r(A) = \min(|A|, k)$  which is a non-decreasing concave function applied to a modular function.
- Closure function

$$\text{span}(A) = \begin{cases} A & \text{if } |A| < k, \\ E & \text{if } |A| \geq k, \end{cases} \quad (4.74)$$

- A “free” matroid sets  $k = |E|$ , so everything is independent.

# Linear (or Matric) Matroid

- Let  $\mathbf{X}$  be an  $n \times m$  matrix and  $E = \{1, \dots, m\}$



# Linear (or Matric) Matroid

- Let  $\mathbf{X}$  be an  $n \times m$  matrix and  $E = \{1, \dots, m\}$
- Let  $\mathcal{I}$  consists of subsets of  $E$  such that if  $A \in \mathcal{I}$ , and  $A = \{a_1, a_2, \dots, a_k\}$  then the vectors  $x_{a_1}, x_{a_2}, \dots, x_{a_k}$  are linearly independent.

# Linear (or Matric) Matroid

- Let  $\mathbf{X}$  be an  $n \times m$  matrix and  $E = \{1, \dots, m\}$
- Let  $\mathcal{I}$  consists of subsets of  $E$  such that if  $A \in \mathcal{I}$ , and  $A = \{a_1, a_2, \dots, a_k\}$  then the vectors  $x_{a_1}, x_{a_2}, \dots, x_{a_k}$  are linearly independent.
- the rank function is just the rank of the space spanned by the corresponding set of vectors.

# Linear (or Matric) Matroid

- Let  $\mathbf{X}$  be an  $n \times m$  matrix and  $E = \{1, \dots, m\}$
- Let  $\mathcal{I}$  consists of subsets of  $E$  such that if  $A \in \mathcal{I}$ , and  $A = \{a_1, a_2, \dots, a_k\}$  then the vectors  $x_{a_1}, x_{a_2}, \dots, x_{a_k}$  are linearly independent.
- the rank function is just the rank of the space spanned by the corresponding set of vectors.
- rank is submodular, it is intuitive that it satisfies the diminishing returns property (a given vector can only become linearly dependent in a greater context, thereby no longer contributing to rank).

# Linear (or Matric) Matroid

- Let  $\mathbf{X}$  be an  $n \times m$  matrix and  $E = \{1, \dots, m\}$
- Let  $\mathcal{I}$  consists of subsets of  $E$  such that if  $A \in \mathcal{I}$ , and  $A = \{a_1, a_2, \dots, a_k\}$  then the vectors  $x_{a_1}, x_{a_2}, \dots, x_{a_k}$  are linearly independent.
- the rank function is just the rank of the space spanned by the corresponding set of vectors.
- rank is submodular, it is intuitive that it satisfies the diminishing returns property (a given vector can only become linearly dependent in a greater context, thereby no longer contributing to rank).
- Called both linear matroids and matric matroids.

# Cycle Matroid of a graph: Graphic Matroids

- Let  $G = (V, E)$  be a graph. Consider  $(E, \mathcal{I})$  where the edges of the graph  $E$  are the ground set and  $A \in \mathcal{I}$  if the edge-induced graph  $G(V, A)$  by  $A$  does not contain any cycle.

# Cycle Matroid of a graph: Graphic Matroids

- Let  $G = (V, E)$  be a graph. Consider  $(E, \mathcal{I})$  where the edges of the graph  $E$  are the ground set and  $A \in \mathcal{I}$  if the edge-induced graph  $G(V, A)$  by  $A$  does not contain any cycle.
- Then  $M = (E, \mathcal{I})$  is a matroid.

# Cycle Matroid of a graph: Graphic Matroids

- Let  $G = (V, E)$  be a graph. Consider  $(E, \mathcal{I})$  where the edges of the graph  $E$  are the ground set and  $A \in \mathcal{I}$  if the edge-induced graph  $G(V, A)$  by  $A$  does not contain any cycle.
- Then  $M = (E, \mathcal{I})$  is a matroid.
- $\mathcal{I}$  contains all forests.

# Cycle Matroid of a graph: Graphic Matroids

- Let  $G = (V, E)$  be a graph. Consider  $(E, \mathcal{I})$  where the edges of the graph  $E$  are the ground set and  $A \in \mathcal{I}$  if the edge-induced graph  $G(V, A)$  by  $A$  does not contain any cycle.
- Then  $M = (E, \mathcal{I})$  is a matroid.
- $\mathcal{I}$  contains all forests.
- Bases are spanning forests (spanning trees if  $G$  is connected).



# Cycle Matroid of a graph: Graphic Matroids


- Let  $G = (V, E)$  be a graph. Consider  $(E, \mathcal{I})$  where the edges of the graph  $E$  are the ground set and  $A \in \mathcal{I}$  if the edge-induced graph  $G(V, A)$  by  $A$  does not contain any cycle.
- Then  $M = (E, \mathcal{I})$  is a matroid.
- $\mathcal{I}$  contains all forests.
- Bases are spanning forests (spanning trees if  $G$  is connected).
- Rank function  $r(A)$  is the size of the largest spanning forest contained in  $G(V, A)$ .

# Cycle Matroid of a graph: Graphic Matroids

- Let  $G = (V, E)$  be a graph. Consider  $(E, \mathcal{I})$  where the edges of the graph  $E$  are the ground set and  $A \in \mathcal{I}$  if the edge-induced graph  $G(V, A)$  by  $A$  does not contain any cycle.
- Then  $M = (E, \mathcal{I})$  is a matroid.
- $\mathcal{I}$  contains all forests.
- Bases are spanning forests (spanning trees if  $G$  is connected).
- Rank function  $r(A)$  is the size of the largest spanning forest contained in  $G(V, A)$ .
- Closure function adds all edges between the vertices adjacent to any edge in  $A$ . Closure of a spanning forest is  $G$ .


# Example: graphic matroid

- A graph defines a matroid on edge sets, independent sets are those without a cycle.




# Example: graphic matroid

- A graph defines a matroid on edge sets, independent sets are those without a cycle.




# Example: graphic matroid

- A graph defines a matroid on edge sets, independent sets are those without a cycle.




# Example: graphic matroid

- A graph defines a matroid on edge sets, independent sets are those without a cycle.




# Example: graphic matroid

- A graph defines a matroid on edge sets, independent sets are those without a cycle.



# Example: graphic matroid

- A graph defines a matroid on edge sets, independent sets are those without a cycle.





# Partition Matroid

- Let  $V$  be our ground set.

# Partition Matroid

- Let  $V$  be our ground set.
- Let  $V = V_1 \cup V_2 \cup \dots \cup V_\ell$  be a partition of  $V$  into blocks or disjoint sets (disjoint union). Define a set of subsets of  $V$  as

$$\mathcal{I} = \{X \subseteq V : |X \cap V_i| \leq k_i \text{ for all } i = 1, \dots, \ell\}. \quad (4.75)$$

where  $k_1, \dots, k_\ell$  are fixed parameters,  $k_i \geq 0$ . Then  $M = (V, \mathcal{I})$  is a matroid.

# Partition Matroid

- Let  $V$  be our ground set.
- Let  $V = V_1 \cup V_2 \cup \dots \cup V_\ell$  be a partition of  $V$  into blocks or disjoint sets (disjoint union). Define a set of subsets of  $V$  as

$$\mathcal{I} = \{X \subseteq V : |X \cap V_i| \leq k_i \text{ for all } i = 1, \dots, \ell\}. \quad (4.75)$$

where  $k_1, \dots, k_\ell$  are fixed parameters,  $k_i \geq 0$ . Then  $M = (V, \mathcal{I})$  is a matroid.

- Note that a  $k$ -uniform matroid is a trivial example of a partition matroid with  $\ell = 1$ ,  $V_1 = V$ , and  $k_1 = k$ .

# Partition Matroid

- Let  $V$  be our ground set.
- Let  $V = V_1 \cup V_2 \cup \dots \cup V_\ell$  be a partition of  $V$  into blocks or disjoint sets (disjoint union). Define a set of subsets of  $V$  as

$$\mathcal{I} = \{X \subseteq V : |X \cap V_i| \leq k_i \text{ for all } i = 1, \dots, \ell\}. \quad (4.75)$$

where  $k_1, \dots, k_\ell$  are fixed parameters,  $k_i \geq 0$ . Then  $M = (V, \mathcal{I})$  is a matroid.

- Note that a  $k$ -uniform matroid is a trivial example of a partition matroid with  $\ell = 1$ ,  $V_1 = V$ , and  $k_1 = k$ .
- We'll show that property (I3') in Def 4.5.6 holds. If  $X, Y \in \mathcal{I}$  with  $|Y| > |X|$ , then there must be at least one  $i$  with  $|Y \cap V_i| > |X \cap V_i|$ . Therefore, adding one element  $e \in V_i \cap (Y \setminus X)$  to  $X$  won't break independence.

# Partition Matroid

Ground set of objects,  $V = \left\{ \right.$



$\left. \right\}$

# Partition Matroid

Partition of  $V$  into six blocks,  $V_1, V_2, \dots, V_6$



# Partition Matroid

Limit associated with each block,  $\{k_1, k_2, \dots, k_6\}$



# Partition Matroid

Independent subset but not maximally independent.





# Partition Matroid

Maximally independent subset, what is called a **base**.



# Partition Matroid

Not independent since over limit in set six.



# Matroids - rank

## Lemma 4.7.1

*The rank function  $r : 2^E \rightarrow \mathbb{Z}_+$  of a matroid is submodular, that is*

$$r(A) + r(B) \geq r(A \cup B) + r(A \cap B)$$

# Matroids - rank

## Lemma 4.7.1

*The rank function  $r : 2^E \rightarrow \mathbb{Z}_+$  of a matroid is submodular, that is*  

$$r(A) + r(B) \geq r(A \cup B) + r(A \cap B)$$

## Proof.

- ① Let  $X \in \mathcal{I}$  be an inclusionwise maximal set with  $X \subseteq A \cap B$



# Matroids - rank

## Lemma 4.7.1

The rank function  $r : 2^E \rightarrow \mathbb{Z}_+$  of a matroid is submodular, that is

$$r(A) + r(B) \geq r(A \cup B) + r(A \cap B)$$

### Proof.

- ① Let  $X \in \mathcal{I}$  be an inclusionwise maximal set with  $X \subseteq A \cap B$
- ② Let  $Y \in \mathcal{I}$  be inclusionwise maximal set with  $X \subseteq Y \subseteq A \cup B$ . (We can find such a  $Y \supseteq X$  because, starting from  $X \subseteq A \cup B$ , and since  $|Y| \geq |X|$ , we can choose a  $y \in Y \subseteq A \cup B$  such that  $X + y \in \mathcal{I}$  but since  $y \in A \cup B$ , also  $X + y \in A \cup B$ . We can keep doing this while  $|Y| > |X|$  since this is a matroid.)

# Matroids - rank

## Lemma 4.7.1

*The rank function  $r : 2^E \rightarrow \mathbb{Z}_+$  of a matroid is submodular, that is*

$$r(A) + r(B) \geq r(A \cup B) + r(A \cap B)$$

### Proof.

- ① Let  $X \in \mathcal{I}$  be an inclusionwise maximal set with  $X \subseteq A \cap B$
- ② Let  $Y \in \mathcal{I}$  be inclusionwise maximal set with  $X \subseteq Y \subseteq A \cup B$ .
- ③ Since  $M$  is a matroid, we know that  $r(A \cap B) = r(X) = |X|$ , and  $r(A \cup B) = r(Y) = |Y|$ . Also, for any  $U \in \mathcal{I}$ ,  $r(A) \geq |A \cap U|$ .



# Matroids - rank

## Lemma 4.7.1

*The rank function  $r : 2^E \rightarrow \mathbb{Z}_+$  of a matroid is submodular, that is*

$$r(A) + r(B) \geq r(A \cup B) + r(A \cap B)$$

### Proof.

- ① Let  $X \in \mathcal{I}$  be an inclusionwise maximal set with  $X \subseteq A \cap B$
- ② Let  $Y \in \mathcal{I}$  be inclusionwise maximal set with  $X \subseteq Y \subseteq A \cup B$ .
- ③ Since  $M$  is a matroid, we know that  $r(A \cap B) = r(X) = |X|$ , and  $r(A \cup B) = r(Y) = |Y|$ . Also, for any  $U \in \mathcal{I}$ ,  $r(A) \geq |A \cap U|$ .
- ④ Then we have

$$r(A) + r(B) \tag{4.76}$$



# Matroids - rank

## Lemma 4.7.1

*The rank function  $r : 2^E \rightarrow \mathbb{Z}_+$  of a matroid is submodular, that is*

$$r(A) + r(B) \geq r(A \cup B) + r(A \cap B)$$

### Proof.

- ① Let  $X \in \mathcal{I}$  be an inclusionwise maximal set with  $X \subseteq A \cap B$
- ② Let  $Y \in \mathcal{I}$  be inclusionwise maximal set with  $X \subseteq Y \subseteq A \cup B$ .
- ③ Since  $M$  is a matroid, we know that  $r(A \cap B) = r(X) = |X|$ , and  $r(A \cup B) = r(Y) = |Y|$ . Also, for any  $U \in \mathcal{I}$ ,  $r(A) \geq |A \cap U|$ .
- ④ Then we have

$$r(A) + r(B) \geq |Y \cap A| + |Y \cap B| \tag{4.76}$$





# Matroids - rank

## Lemma 4.7.1

*The rank function  $r : 2^E \rightarrow \mathbb{Z}_+$  of a matroid is submodular, that is*

$$r(A) + r(B) \geq r(A \cup B) + r(A \cap B)$$

### Proof.

- ① Let  $X \in \mathcal{I}$  be an inclusionwise maximal set with  $X \subseteq A \cap B$
- ② Let  $Y \in \mathcal{I}$  be inclusionwise maximal set with  $X \subseteq Y \subseteq A \cup B$ .
- ③ Since  $M$  is a matroid, we know that  $r(A \cap B) = r(X) = |X|$ , and  $r(A \cup B) = r(Y) = |Y|$ . Also, for any  $U \in \mathcal{I}$ ,  $r(A) \geq |A \cap U|$ .
- ④ Then we have

$$r(A) + r(B) \geq |Y \cap A| + |Y \cap B| \tag{4.76}$$

$$= |Y \cap (A \cap B)| + |Y \cap (A \cup B)| \tag{4.77}$$



# Matroids - rank

## Lemma 4.7.1

The rank function  $r : 2^E \rightarrow \mathbb{Z}_+$  of a matroid is submodular, that is

$$r(A) + r(B) \geq r(A \cup B) + r(A \cap B)$$

### Proof.

- ① Let  $X \in \mathcal{I}$  be an inclusionwise maximal set with  $X \subseteq A \cap B$
- ② Let  $Y \in \mathcal{I}$  be inclusionwise maximal set with  $X \subseteq Y \subseteq A \cup B$ .
- ③ Since  $M$  is a matroid, we know that  $r(A \cap B) = r(X) = |X|$ , and  $r(A \cup B) = r(Y) = |Y|$ . Also, for any  $U \in \mathcal{I}$ ,  $r(A) \geq |A \cap U|$ .
- ④ Then we have

$$r(A) + r(B) \geq |Y \cap A| + |Y \cap B| \tag{4.76}$$

$$= |Y \cap (A \cap B)| + |Y \cap (A \cup B)| \tag{4.77}$$

$$\geq |X| + |Y| = r(A \cap B) + r(A \cup B) \tag{4.78}$$



# Matroids

In fact, we can use the rank of a matroid for its definition.

## Theorem 4.7.2 (Matroid from rank)

*Let  $E$  be a set and let  $r : 2^E \rightarrow \mathbb{Z}_+$  be a function. Then  $r(\cdot)$  defines a matroid with  $r$  being its rank function if and only if for all  $A, B \subseteq E$ :*

- (R1)  $\forall A \subseteq E \quad 0 \leq r(A) \leq |A|$  (non-negative cardinality bounded)*
- (R2)  $r(A) \leq r(B)$  whenever  $A \subseteq B \subseteq E$  (monotone non-decreasing)*
- (R3)  $r(A \cup B) + r(A \cap B) \leq r(A) + r(B)$  for all  $A, B \subseteq E$  (submodular)*

- So submodularity and non-negative monotone non-decreasing, and unit increase is necessary and sufficient to define the matroid.
- Given above, unit increment (if  $r(A) = k$ , then either  $r(A \cup \{v\}) = k$  or  $r(A \cup \{v\}) = k + 1$ ) holds.
- A matroid is sometimes given as  $(E, r)$  where  $E$  is ground set and  $r$  is rank function.

# Matroids

In fact, we can use the rank of a matroid for its definition.

## Theorem 4.7.2 (Matroid from rank)

Let  $E$  be a set and let  $r : 2^E \rightarrow \mathbb{Z}_+$  be a function. Then  $r(\cdot)$  defines a matroid with  $r$  being its rank function if and only if for all  $A, B \subseteq E$ :

- (R1)  $\forall A \subseteq E \quad 0 \leq r(A) \leq |A|$  (non-negative cardinality bounded)
- (R2)  $r(A) \leq r(B)$  whenever  $A \subseteq B \subseteq E$  (monotone non-decreasing)
- (R3)  $r(A \cup B) + r(A \cap B) \leq r(A) + r(B)$  for all  $A, B \subseteq E$  (submodular)

- From above,  $r(\emptyset) = 0$ . Let  $v \notin A$ , then by monotonicity and submodularity,  $r(A) \leq r(A \cup \{v\}) \leq r(A) + r(\{v\})$  which gives only two possible values to  $r(A \cup \{v\})$ .

# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank).

- Given a matroid  $M = (E, \mathcal{I})$ , we see its rank function as defined in Eq. 4.71 satisfies (R1), (R2), and, as we saw in Lemma 4.7.1, (R3) too.

...

# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank).

- Given a matroid  $M = (E, \mathcal{I})$ , we see its rank function as defined in Eq. 4.71 satisfies (R1), (R2), and, as we saw in Lemma 4.7.1, (R3) too.
- Next, assume we have (R1), (R2), and (R3). Define  $\mathcal{I} = \{X \subseteq E : r(X) = |X|\}$ . We will show that  $(E, \mathcal{I})$  is a matroid.

...

# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank).

- Given a matroid  $M = (E, \mathcal{I})$ , we see its rank function as defined in Eq. 4.71 satisfies (R1), (R2), and, as we saw in Lemma 4.7.1, (R3) too.
- Next, assume we have (R1), (R2), and (R3). Define  $\mathcal{I} = \{X \subseteq E : r(X) = |X|\}$ . We will show that  $(E, \mathcal{I})$  is a matroid.
- First,  $\emptyset \in \mathcal{I}$ .

...

# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank).

- Given a matroid  $M = (E, \mathcal{I})$ , we see its rank function as defined in Eq. 4.71 satisfies (R1), (R2), and, as we saw in Lemma 4.7.1, (R3) too.
- Next, assume we have (R1), (R2), and (R3). Define  $\mathcal{I} = \{X \subseteq E : r(X) = |X|\}$ . We will show that  $(E, \mathcal{I})$  is a matroid.
- First,  $\emptyset \in \mathcal{I}$ .
- Also, if  $Y \in \mathcal{I}$  and  $X \subseteq Y$  then by submodularity,

...



# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank).

- Given a matroid  $M = (E, \mathcal{I})$ , we see its rank function as defined in Eq. 4.71 satisfies (R1), (R2), and, as we saw in Lemma 4.7.1, (R3) too.
- Next, assume we have (R1), (R2), and (R3). Define  $\mathcal{I} = \{X \subseteq E : r(X) = |X|\}$ . We will show that  $(E, \mathcal{I})$  is a matroid.
- First,  $\emptyset \in \mathcal{I}$ .
- Also, if  $Y \in \mathcal{I}$  and  $X \subseteq Y$  then by submodularity,

$$r(X) \geq r(Y) - r(Y \setminus X) \quad (4.79)$$

...

# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank).

- Given a matroid  $M = (E, \mathcal{I})$ , we see its rank function as defined in Eq. 4.71 satisfies (R1), (R2), and, as we saw in Lemma 4.7.1, (R3) too.
- Next, assume we have (R1), (R2), and (R3). Define  $\mathcal{I} = \{X \subseteq E : r(X) = |X|\}$ . We will show that  $(E, \mathcal{I})$  is a matroid.
- First,  $\emptyset \in \mathcal{I}$ .
- Also, if  $Y \in \mathcal{I}$  and  $X \subseteq Y$  then by submodularity,

$$r(X) \geq r(Y) - r(Y \setminus X) - r(\emptyset) \quad (4.79)$$

...

# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank).

- Given a matroid  $M = (E, \mathcal{I})$ , we see its rank function as defined in Eq. 4.71 satisfies (R1), (R2), and, as we saw in Lemma 4.7.1, (R3) too.
- Next, assume we have (R1), (R2), and (R3). Define  $\mathcal{I} = \{X \subseteq E : r(X) = |X|\}$ . We will show that  $(E, \mathcal{I})$  is a matroid.
- First,  $\emptyset \in \mathcal{I}$ .
- Also, if  $Y \in \mathcal{I}$  and  $X \subseteq Y$  then by submodularity,

$$r(X) \geq r(Y) - r(Y \setminus X) - r(\emptyset) \tag{4.79}$$

$$\geq |Y| - |Y \setminus X| \tag{4.80}$$

...

# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank).

- Given a matroid  $M = (E, \mathcal{I})$ , we see its rank function as defined in Eq. 4.71 satisfies (R1), (R2), and, as we saw in Lemma 4.7.1, (R3) too.
- Next, assume we have (R1), (R2), and (R3). Define  $\mathcal{I} = \{X \subseteq E : r(X) = |X|\}$ . We will show that  $(E, \mathcal{I})$  is a matroid.
- First,  $\emptyset \in \mathcal{I}$ .
- Also, if  $Y \in \mathcal{I}$  and  $X \subseteq Y$  then by submodularity,

$$r(X) \geq r(Y) - r(Y \setminus X) - r(\emptyset) \tag{4.79}$$

$$\geq |Y| - |Y \setminus X| \tag{4.80}$$

$$= |X| \tag{4.81}$$

...

# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank).

- Given a matroid  $M = (E, \mathcal{I})$ , we see its rank function as defined in Eq. 4.71 satisfies (R1), (R2), and, as we saw in Lemma 4.7.1, (R3) too.
- Next, assume we have (R1), (R2), and (R3). Define  $\mathcal{I} = \{X \subseteq E : r(X) = |X|\}$ . We will show that  $(E, \mathcal{I})$  is a matroid.
- First,  $\emptyset \in \mathcal{I}$ .
- Also, if  $Y \in \mathcal{I}$  and  $X \subseteq Y$  then by submodularity,

$$r(X) \geq r(Y) - r(Y \setminus X) - r(\emptyset) \tag{4.79}$$

$$\geq |Y| - |Y \setminus X| \tag{4.80}$$

$$= |X| \tag{4.81}$$

implying  $r(X) = |X|$ , and thus  $X \in \mathcal{I}$ .

...

# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank) cont.

- Let  $A, B \in \mathcal{I}$ , with  $|A| < |B|$ , so  $r(A) = |A| < r(B) = |B|$ . Let  $B \setminus A = \{b_1, b_2, \dots, b_k\}$  (note  $k \leq |B|$ ).



# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank) cont.

- Let  $A, B \in \mathcal{I}$ , with  $|A| < |B|$ , so  $r(A) = |A| < r(B) = |B|$ . Let  $B \setminus A = \{b_1, b_2, \dots, b_k\}$  (note  $k \leq |B|$ ).
- Suppose, to the contrary, that  $\forall b \in B \setminus A, r(A + b) \notin \mathcal{I}$ , which means for all such  $b$ ,  $r(A + b) = r(A) = |A|$ . Then



# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank) cont.

- Let  $A, B \in \mathcal{I}$ , with  $|A| < |B|$ , so  $r(A) = |A| < r(B) = |B|$ . Let  $B \setminus A = \{b_1, b_2, \dots, b_k\}$  (note  $k \leq |B|$ ).
- Suppose, to the contrary, that  $\forall b \in B \setminus A, r(A + b) \notin \mathcal{I}$ , which means for all such  $b$ ,  $r(A + b) = r(A) = |A|$ . Then

$$r(B) \leq r(A \cup B) \tag{4.82}$$





# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank) cont.

- Let  $A, B \in \mathcal{I}$ , with  $|A| < |B|$ , so  $r(A) = |A| < r(B) = |B|$ . Let  $B \setminus A = \{b_1, b_2, \dots, b_k\}$  (note  $k \leq |B|$ ).
- Suppose, to the contrary, that  $\forall b \in B \setminus A, r(A + b) \notin \mathcal{I}$ , which means for all such  $b$ ,  $r(A + b) = r(A) = |A|$ . Then

$$r(B) \leq r(A \cup B) \tag{4.82}$$

$$\leq r(A \cup (B \setminus \{b_1\})) + r(A \cup \{b_1\}) - r(A) \tag{4.83}$$



# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank) cont.

- Let  $A, B \in \mathcal{I}$ , with  $|A| < |B|$ , so  $r(A) = |A| < r(B) = |B|$ . Let  $B \setminus A = \{b_1, b_2, \dots, b_k\}$  (note  $k \leq |B|$ ).
- Suppose, to the contrary, that  $\forall b \in B \setminus A, r(A + b) \notin \mathcal{I}$ , which means for all such  $b$ ,  $r(A + b) = r(A) = |A|$ . Then

$$r(B) \leq r(A \cup B) \tag{4.82}$$

$$\leq r(A \cup (B \setminus \{b_1\})) + r(A \cup \{b_1\}) - r(A) \tag{4.83}$$

$$= r(A \cup (B \setminus \{b_1\})) \tag{4.84}$$



# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank) cont.

- Let  $A, B \in \mathcal{I}$ , with  $|A| < |B|$ , so  $r(A) = |A| < r(B) = |B|$ . Let  $B \setminus A = \{b_1, b_2, \dots, b_k\}$  (note  $k \leq |B|$ ).
- Suppose, to the contrary, that  $\forall b \in B \setminus A, r(A + b) \notin \mathcal{I}$ , which means for all such  $b$ ,  $r(A + b) = r(A) = |A|$ . Then

$$r(B) \leq r(A \cup B) \tag{4.82}$$

$$\leq r(A \cup (B \setminus \{b_1\})) + r(A \cup \{b_1\}) - r(A) \tag{4.83}$$

$$= r(A \cup (B \setminus \{b_1\})) \tag{4.84}$$

$$\leq r(A \cup (B \setminus \{b_1, b_2\})) + r(A \cup \{b_2\}) - r(A) \tag{4.85}$$



# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank) cont.

- Let  $A, B \in \mathcal{I}$ , with  $|A| < |B|$ , so  $r(A) = |A| < r(B) = |B|$ . Let  $B \setminus A = \{b_1, b_2, \dots, b_k\}$  (note  $k \leq |B|$ ).
- Suppose, to the contrary, that  $\forall b \in B \setminus A, r(A + b) \notin \mathcal{I}$ , which means for all such  $b, r(A + b) = r(A) = |A|$ . Then

$$r(B) \leq r(A \cup B) \tag{4.82}$$

$$\leq r(A \cup (B \setminus \{b_1\})) + r(A \cup \{b_1\}) - r(A) \tag{4.83}$$

$$= r(A \cup (B \setminus \{b_1\})) \tag{4.84}$$

$$\leq r(A \cup (B \setminus \{b_1, b_2\})) + r(A \cup \{b_2\}) - r(A) \tag{4.85}$$

$$= r(A \cup (B \setminus \{b_1, b_2\})) \tag{4.86}$$



# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank) cont.

- Let  $A, B \in \mathcal{I}$ , with  $|A| < |B|$ , so  $r(A) = |A| < r(B) = |B|$ . Let  $B \setminus A = \{b_1, b_2, \dots, b_k\}$  (note  $k \leq |B|$ ).
- Suppose, to the contrary, that  $\forall b \in B \setminus A, r(A + b) \notin \mathcal{I}$ , which means for all such  $b, r(A + b) = r(A) = |A|$ . Then

$$r(B) \leq r(A \cup B) \tag{4.82}$$

$$\leq r(A \cup (B \setminus \{b_1\})) + r(A \cup \{b_1\}) - r(A) \tag{4.83}$$

$$= r(A \cup (B \setminus \{b_1\})) \tag{4.84}$$

$$\leq r(A \cup (B \setminus \{b_1, b_2\})) + r(A \cup \{b_2\}) - r(A) \tag{4.85}$$

$$= r(A \cup (B \setminus \{b_1, b_2\})) \tag{4.86}$$

$$\leq \dots \leq r(A) = |A| < |B| \tag{4.87}$$



# Matroids from rank

## Proof of Theorem 4.7.2 (matroid from rank) cont.

- Let  $A, B \in \mathcal{I}$ , with  $|A| < |B|$ , so  $r(A) = |A| < r(B) = |B|$ . Let  $B \setminus A = \{b_1, b_2, \dots, b_k\}$  (note  $k \leq |B|$ ).
- Suppose, to the contrary, that  $\forall b \in B \setminus A, r(A + b) \notin \mathcal{I}$ , which means for all such  $b$ ,  $r(A + b) = r(A) = |A|$ . Then

$$r(B) \leq r(A \cup B) \tag{4.82}$$

$$\leq r(A \cup (B \setminus \{b_1\})) + r(A \cup \{b_1\}) - r(A) \tag{4.83}$$

$$= r(A \cup (B \setminus \{b_1\})) \tag{4.84}$$

$$\leq r(A \cup (B \setminus \{b_1, b_2\})) + r(A \cup \{b_2\}) - r(A) \tag{4.85}$$

$$= r(A \cup (B \setminus \{b_1, b_2\})) \tag{4.86}$$

$$\leq \dots \leq r(A) = |A| < |B| \tag{4.87}$$

giving a contradiction since  $B \in \mathcal{I}$ .



# Matroids from rank II

Another way of using function  $r$  to define a matroid.

## Theorem 4.7.3 (Matroid from rank II)

*Let  $E$  be a finite set and let  $r : 2^E \rightarrow \mathbb{Z}_+$  be a function. Then  $r(\cdot)$  defines a matroid with  $r$  being its rank function if and only if for all  $A \subseteq E$ , and  $x, y \in E$ :*

- (R1')  $r(\emptyset) = 0$ ;
- (R2')  $r(X) \leq r(X \cup \{y\}) \leq r(X) + 1$ ;
- (R3') *If  $r(X \cup \{x\}) = r(X \cup \{y\}) = r(X)$ , then  $r(X \cup \{x, y\}) = r(X)$ .*

# Matroid and Rank

- Thus, we can define a matroid as  $M = (V, r)$  where  $r$  satisfies matroid rank axioms.
- Example: 2-partition matroid rank function: Given natural numbers  $a, b \in \mathbb{Z}_+$  with  $a > b$ , and any set  $R \subseteq V$  with  $|R| = a$ , two-block partition  $V = (R, \bar{R})$ , define:

$$r(A) = \min(|A \cap R|, b) + \min(|A \cap \bar{R}|, |\bar{R}|) \quad (4.88)$$

$$= \min(|A \cap R|, b) + |A \cap \bar{R}| \quad (4.89)$$

- Example: **Truncated matroid rank** function.

$$f_R(A) = \min \{r(A), a\} \quad (4.90)$$

$$= \min \{|A|, b + |A \cap \bar{R}|, a\} \quad (4.91)$$

- Defines a matroid  $M = (V, f_R) = (V, \mathcal{I})$  (Goemans et. al.) with

$$\mathcal{I} = \{I \subseteq V : |I| \leq a \text{ and } |I \cap R| \leq b\}, \quad (4.92)$$

useful for showing hardness of constrained submodular minimization.



# Summarizing: Many ways to define a Matroid

Summarizing what we've so far seen, we saw that it is possible to uniquely define a matroid based on any of:

- Independence (define the independent sets).

# Summarizing: Many ways to define a Matroid

Summarizing what we've so far seen, we saw that it is possible to uniquely define a matroid based on any of:

- Independence (define the independent sets).
- Base axioms (exchangeability)

# Summarizing: Many ways to define a Matroid

Summarizing what we've so far seen, we saw that it is possible to uniquely define a matroid based on any of:

- Independence (define the independent sets).
- Base axioms (exchangeability)
- Circuit axioms

# Summarizing: Many ways to define a Matroid

Summarizing what we've so far seen, we saw that it is possible to uniquely define a matroid based on any of:

- Independence (define the independent sets).
- Base axioms (exchangeability)
- Circuit axioms
- Closure axioms (we didn't see this, but it is possible)

# Summarizing: Many ways to define a Matroid

Summarizing what we've so far seen, we saw that it is possible to uniquely define a matroid based on any of:

- Independence (define the independent sets).
- Base axioms (exchangeability)
- Circuit axioms
- Closure axioms (we didn't see this, but it is possible)
- Rank axioms (normalized, monotone, cardinality bounded, submodular)

# Maximization problems for matroids

- Given a matroid  $M = (E, \mathcal{I})$  and a modular cost function  $c : E \rightarrow \mathbb{R}$ , the task is to find an  $X \in \mathcal{I}$  such that  $c(X) = \sum_{x \in X} c(x)$  is maximum.
- This seems remarkably similar to the max spanning tree problem.

# Minimization problems for matroids

- Given a matroid  $M = (E, \mathcal{I})$  and a modular cost function  $c : E \rightarrow \mathbb{R}$ , the task is to find a basis  $B \in \mathcal{B}$  such that  $c(B)$  is minimized.
- This sounds like a set cover problem (find the minimum cost covering set of sets).

# Partition Matroid

- What is the partition matroid's rank function?



# Partition Matroid

- What is the partition matroid's rank function?
- A partition matroid's rank function:

$$r(A) = \sum_{i=1}^{\ell} \min(|A \cap V_i|, k_i) \quad (4.93)$$

which we also immediately see is submodular using properties we spoke about last week. That is:

# Partition Matroid

- What is the partition matroid's rank function?
- A partition matroid's rank function:

$$r(A) = \sum_{i=1}^{\ell} \min(|A \cap V_i|, k_i) \quad (4.93)$$

which we also immediately see is submodular using properties we spoke about last week. That is:

- ①  $|A \cap V_i|$  is submodular (even modular) in  $A$

# Partition Matroid

- What is the partition matroid's rank function?
- A partition matroid's rank function:

$$r(A) = \sum_{i=1}^{\ell} \min(|A \cap V_i|, k_i) \quad (4.93)$$

which we also immediately see is submodular using properties we spoke about last week. That is:

- 1  $|A \cap V_i|$  is submodular (even modular) in  $A$
- 2  $\min(\text{submodular}(A), k_i)$  is submodular in  $A$  since  $|A \cap V_i|$  is monotone.

# Partition Matroid

- What is the partition matroid's rank function?
- A partition matroid's rank function:

$$r(A) = \sum_{i=1}^{\ell} \min(|A \cap V_i|, k_i) \quad (4.93)$$

which we also immediately see is submodular using properties we spoke about last week. That is:

- 1  $|A \cap V_i|$  is submodular (even modular) in  $A$
- 2  $\min(\text{submodular}(A), k_i)$  is submodular in  $A$  since  $|A \cap V_i|$  is monotone.
- 3 sums of submodular functions are submodular.

# Partition Matroid

- What is the partition matroid's rank function?
- A partition matroid's rank function:

$$r(A) = \sum_{i=1}^{\ell} \min(|A \cap V_i|, k_i) \quad (4.93)$$

which we also immediately see is submodular using properties we spoke about last week. That is:


- 1  $|A \cap V_i|$  is submodular (even modular) in  $A$
  - 2  $\min(\text{submodular}(A), k_i)$  is submodular in  $A$  since  $|A \cap V_i|$  is monotone.
  - 3 sums of submodular functions are submodular.
- $r(A)$  is also non-negative integral monotone non-decreasing, so it defines a matroid (the partition matroid).

# Partition Matroid, rank as matching

- A partition matroid can be viewed using a bipartite graph.
- Letting  $V$  denote the ground set, and  $V_1, V_2, \dots$  the partition, the graph is  $G = (V, I, E)$  where  $V$  is the ground set,  $I$  is a set of “indices”, and  $E$  is the set of edges.
- $I = (I_1, I_2, \dots, I_\ell)$  is a set of  $k = \sum_{i=1}^{\ell} k_i$  nodes, grouped into  $\ell$  clusters, where there are  $k_i$  nodes in the  $i^{\text{th}}$  group  $I_i$ .
- $(v, i) \in E(G)$  iff  $v \in V_j$  and  $i \in I_j$ .


# Partition Matroid, rank as matching

- Example where  $\ell = 5$ ,  
 $(k_1, k_2, k_3, k_4, k_5) =$   
 $(2, 2, 1, 1, 3).$



# Partition Matroid, rank as matching

- Example where  $\ell = 5$ ,  
 $(k_1, k_2, k_3, k_4, k_5) =$   
 $(2, 2, 1, 1, 3)$ .




- Recall,  $\Gamma : 2^V \rightarrow \mathbb{R}$  as the neighbor function in a bipartite graph, the neighbors of  $X$  is defined as  $\Gamma(X) = \{v \in V(G) \setminus X : E(X, \{v\}) \neq \emptyset\}$ , and recall that  $|\Gamma(X)|$  is submodular.



# Partition Matroid, rank as matching


- Example where  $\ell = 5$ ,  
 $(k_1, k_2, k_3, k_4, k_5) =$   
 $(2, 2, 1, 1, 3)$ .



- Recall,  $\Gamma : 2^V \rightarrow \mathbb{R}$  as the neighbor function in a bipartite graph, the neighbors of  $X$  is defined as  $\Gamma(X) = \{v \in V(G) \setminus X : E(X, \{v\}) \neq \emptyset\}$ , and recall that  $|\Gamma(X)|$  is submodular.
- Here, for  $X \subseteq V$ , we have  $\Gamma(X) = \{i \in I : (v, i) \in E(G) \text{ and } v \in X\}$ .

# Partition Matroid, rank as matching

- Example where  $\ell = 5$ ,  
 $(k_1, k_2, k_3, k_4, k_5) =$   
 $(2, 2, 1, 1, 3)$ .




- Recall,  $\Gamma : 2^V \rightarrow \mathbb{R}$  as the neighbor function in a bipartite graph, the neighbors of  $X$  is defined as  $\Gamma(X) = \{v \in V(G) \setminus X : E(X, \{v\}) \neq \emptyset\}$ , and recall that  $|\Gamma(X)|$  is submodular.
- Here, for  $X \subseteq V$ , we have  $\Gamma(X) = \{i \in I : (v, i) \in E(G) \text{ and } v \in X\}$ .
- For such a constructed bipartite graph, the rank function of a partition matroid is  $r(X) = \sum_{i=1}^{\ell} \min(|X \cap V_i|, k_i) =$  maximum matching involving  $X$ .

# Laminar Matroid

- We can define a matroid with structures richer than just partitions.


# Laminar Matroid

- We can define a matroid with structures richer than just partitions.
- A set system  $(V, \mathcal{F})$  is called a **laminar** family if for any two sets  $A, B \in \mathcal{F}$ , at least one of the three sets  $A \cap B$ ,  $A \setminus B$ , or  $B \setminus A$  is empty.



# Laminar Matroid


- We can define a matroid with structures richer than just partitions.
- A set system  $(V, \mathcal{F})$  is called a **laminar** family if for any two sets  $A, B \in \mathcal{F}$ , at least one of the three sets  $A \cap B$ ,  $A \setminus B$ , or  $B \setminus A$  is empty.



- Family is laminar if it has no two “properly intersecting” members: i.e., intersecting  $A \cap B \neq \emptyset$  and not comparable (one is not contained in the other).

# Laminar Matroid


- We can define a matroid with structures richer than just partitions.
- A set system  $(V, \mathcal{F})$  is called a **laminar** family if for any two sets  $A, B \in \mathcal{F}$ , at least one of the three sets  $A \cap B$ ,  $A \setminus B$ , or  $B \setminus A$  is empty.



- Family is laminar if it has no two “properly intersecting” members: i.e., intersecting  $A \cap B \neq \emptyset$  and not comparable (one is not contained in the other).
- Suppose we have a laminar family  $\mathcal{F}$  of subsets of  $V$  and an integer  $k(A)$  for every set  $A \in \mathcal{F}$ .

# Laminar Matroid

- We can define a matroid with structures richer than just partitions.
- A set system  $(V, \mathcal{F})$  is called a **laminar** family if for any two sets  $A, B \in \mathcal{F}$ , at least one of the three sets  $A \cap B$ ,  $A \setminus B$ , or  $B \setminus A$  is empty.



- Family is laminar if it has no two “properly intersecting” members: i.e., intersecting  $A \cap B \neq \emptyset$  and not comparable (one is not contained in the other).
- Suppose we have a laminar family  $\mathcal{F}$  of subsets of  $V$  and an integer  $k(A)$  for every set  $A \in \mathcal{F}$ .
- Then  $(V, \mathcal{I})$  defines a matroid where

$$\mathcal{I} = \{I \subseteq E : |I \cap A| \leq k(A) \text{ for all } A \in \mathcal{F}\} \quad (4.94)$$

# System of Representatives

- Let  $(V, \mathcal{V})$  be a set system (i.e.,  $\mathcal{V} = (V_k : k \in I)$  where  $\emptyset \subset V_i \subseteq V$  for all  $i$ ).



# System of Representatives

- Let  $(V, \mathcal{V})$  be a set system (i.e.,  $\mathcal{V} = (V_k : k \in I)$  where  $\emptyset \subset V_k \subseteq V$  for all  $k$ ).
- A family  $(v_i : i \in I)$  with  $v_i \in V$  for index set  $I$  is said to be a **system of representatives** of  $\mathcal{V}$  if  $\exists$  a bijection  $\pi : I \rightarrow I$  such that  $v_i \in V_{\pi(i)}$ .  *$v_i$  is the representative of set  $\pi(i)$ , meaning the  $i^{\text{th}}$  representative is meant to represent set  $V_{\pi(i)}$ . Consider the house of representatives,  $v_i = \text{"John Smith"}$ , while  $i = \text{King County}$ .*

# System of Representatives


- Let  $(V, \mathcal{V})$  be a set system (i.e.,  $\mathcal{V} = (V_k : k \in I)$  where  $\emptyset \subset V_i \subseteq V$  for all  $i$ ).
- A family  $(v_i : i \in I)$  with  $v_i \in V$  for index set  $I$  is said to be a **system of representatives** of  $\mathcal{V}$  if  $\exists$  a bijection  $\pi : I \rightarrow I$  such that  $v_i \in V_{\pi(i)}$ .
- In a system of representatives, there is no requirement for the representatives to be distinct. I.e., we could have  $v_1 \in T$ , where  $v_1$  represents both  $V_1$  and  $V_2$ .

# System of Representatives

- Let  $(V, \mathcal{V})$  be a set system (i.e.,  $\mathcal{V} = (V_k : k \in I)$  where  $\emptyset \subset V_i \subseteq V$  for all  $i$ ).
- A family  $(v_i : i \in I)$  with  $v_i \in V$  for index set  $I$  is said to be a **system of representatives** of  $\mathcal{V}$  if  $\exists$  a bijection  $\pi : I \rightarrow I$  such that  $v_i \in V_{\pi(i)}$ .
- In a system of representatives, there is no requirement for the representatives to be distinct. I.e., we could have  $v_1 \in T$ , where  $v_1$  represents both  $V_1$  and  $V_2$ .
- We can view this as a bipartite graph.


# System of Representatives

- We can view this as a bipartite graph. The groups of  $V$  are marked by color tags on the left, and also via right neighbors in the graph.
- Here,  $\ell = 6$ , and  $\mathcal{V} = (V_1, V_2, \dots, V_6)$   
 $= (\{e, f, h\}, \{d, e, g\}, \{b, c, e, h\}, \{a, b, h\}, \{a\}, \{a\})$ .



# System of Representatives


- We can view this as a bipartite graph. The groups of  $V$  are marked by color tags on the left, and also via right neighbors in the graph.
- Here,  $\ell = 6$ , and  $\mathcal{V} = (V_1, V_2, \dots, V_6)$   
 $= (\{e, f, h\}, \{d, e, g\}, \{b, c, e, h\}, \{a, b, h\}, \{a\}, \{a\})$ .



- A system of representatives would make sure that there is a representative for each color group. For example,

# System of Representatives


- We can view this as a bipartite graph. The groups of  $V$  are marked by color tags on the left, and also via right neighbors in the graph.
- Here,  $\ell = 6$ , and  $\mathcal{V} = (V_1, V_2, \dots, V_6)$   
 $= (\{e, f, h\}, \{d, e, g\}, \{b, c, e, h\}, \{a, b, h\}, \{a\}, \{a\})$ .



- A system of representatives would make sure that there is a representative for each color group. For example,
- The representatives are shown as colors on the left.

# System of Representatives

- We can view this as a bipartite graph. The groups of  $V$  are marked by color tags on the left, and also via right neighbors in the graph.
- Here,  $\ell = 6$ , and  $\mathcal{V} = (V_1, V_2, \dots, V_6)$   
 $= (\{e, f, h\}, \{d, e, g\}, \{b, c, e, h\}, \{a, b, h\}, \{a\}, \{a\})$ .



- A system of representatives would make sure that there is a representative for each color group. For example,
- The representatives are shown as colors on the left.
- Here, the set of representatives is not distinct. In fact, due to the red and pink group, a distinct group of representatives is impossible (since there is only one common choice to represent both color groups).

# System of Distinct Representatives

- Let  $(V, \mathcal{V})$  be a set system (i.e.,  $\mathcal{V} = (V_k : k \in I)$  where  $V_i \subseteq V$  for all  $i$ ).



# System of Distinct Representatives

- Let  $(V, \mathcal{V})$  be a set system (i.e.,  $\mathcal{V} = (V_k : k \in I)$  where  $V_i \subseteq V$  for all  $i$ ). Hence,  $|I| = |\mathcal{V}|$ .

# System of Distinct Representatives

- Let  $(V, \mathcal{V})$  be a set system (i.e.,  $\mathcal{V} = (V_k : k \in I)$  where  $V_k \subseteq V$  for all  $k$ ). Hence,  $|I| = |\mathcal{V}|$ .
- A family  $(v_i : i \in I)$  with  $v_i \in V$  for index set  $I$  is said to be a **system of distinct representatives** of  $\mathcal{V}$  if  $\exists$  a bijection  $\pi : I \leftrightarrow I$  such that  $v_i \in V_{\pi(i)}$  and  $v_i \neq v_j$  for all  $i \neq j$ .

# System of Distinct Representatives

- Let  $(V, \mathcal{V})$  be a set system (i.e.,  $\mathcal{V} = (V_k : k \in I)$  where  $V_k \subseteq V$  for all  $k$ ). Hence,  $|I| = |\mathcal{V}|$ .
- A family  $(v_i : i \in I)$  with  $v_i \in V$  for index set  $I$  is said to be a **system of distinct representatives** of  $\mathcal{V}$  if  $\exists$  a bijection  $\pi : I \leftrightarrow I$  such that  $v_i \in V_{\pi(i)}$  and  $v_i \neq v_j$  for all  $i \neq j$ .
- In a system of distinct representatives, there **is** a requirement for the representatives to be distinct. Lets re-state (and rename) this as a:

# System of Distinct Representatives

- Let  $(V, \mathcal{V})$  be a set system (i.e.,  $\mathcal{V} = (V_k : k \in I)$  where  $V_i \subseteq V$  for all  $i$ ). Hence,  $|I| = |\mathcal{V}|$ .
- A family  $(v_i : i \in I)$  with  $v_i \in V$  for index set  $I$  is said to be a **system of distinct representatives** of  $\mathcal{V}$  if  $\exists$  a bijection  $\pi : I \leftrightarrow I$  such that  $v_i \in V_{\pi(i)}$  and  $v_i \neq v_j$  for all  $i \neq j$ .
- In a system of distinct representatives, there **is** a requirement for the representatives to be distinct. Lets re-state (and rename) this as a:

## Definition 4.9.1 (transversal)

Given a set system  $(V, \mathcal{V})$  as defined above, a set  $T \subseteq V$  is a **transversal** of  $\mathcal{V}$  if there is a bijection  $\pi : T \leftrightarrow I$  such that

$$x \in V_{\pi(x)} \text{ for all } x \in T \quad (4.95)$$

# System of Distinct Representatives

- Let  $(V, \mathcal{V})$  be a set system (i.e.,  $\mathcal{V} = (V_k : k \in I)$  where  $V_k \subseteq V$  for all  $k$ ). Hence,  $|I| = |\mathcal{V}|$ .
- A family  $(v_i : i \in I)$  with  $v_i \in V$  for index set  $I$  is said to be a **system of distinct representatives** of  $\mathcal{V}$  if  $\exists$  a bijection  $\pi : I \leftrightarrow I$  such that  $v_i \in V_{\pi(i)}$  and  $v_i \neq v_j$  for all  $i \neq j$ .
- In a system of distinct representatives, there **is** a requirement for the representatives to be distinct. Lets re-state (and rename) this as a:

## Definition 4.9.1 (transversal)

Given a set system  $(V, \mathcal{V})$  as defined above, a set  $T \subseteq V$  is a **transversal** of  $\mathcal{V}$  if there is a bijection  $\pi : T \leftrightarrow I$  such that

$$x \in V_{\pi(x)} \text{ for all } x \in T \quad (4.95)$$

- Note that due to it being a bijection, all of  $I$  and  $T$  are “covered” (so this makes things distinct).